arXiv:dg-ga/9604006v1 23 Apr 1996

Rotationally Symmetric p-harmonic maps

Man Chun Leung
Department of Mathematics, National University of Singapore,
Singapore 119260 *matlmc@nus.sg*

April 1996

Abstract

This is a report on our study of a second order ordinary differential equation corresponding to rotationally symmetric p-harmonic maps. We show unique continuation and Liouville's type theorems for positive solutions. Asymptotic properties of the positive solutions are investigated.

KEY WORDS: p-harmonic map, positive solution, rotationally symmetric

1991 AMS MS Classifications: 34C11, 34D05, 58E20

1. Introduction

In this report we study p-harmonic maps between *model* Riemannian manifolds. For $n \ge 2$, let $(S^{n-1}, d\vartheta^2)$ be the unit sphere in $\mathbb{R}^n$ with the induced Riemannian metric $d\vartheta^2$. Let f and g be functions in $C^2([0,\infty))$ which satisfy the following

$$f(0) = g(0) = 0\,, \quad f'(0) = g'(0) = 1 \quad \text{and} \quad f(r), g(r) > 0\,. \tag{1.1}$$

Consider the following model Riemannian manifolds [5]

$$M(f) = ([0,\infty) \times S^{n-1}\,,\ dr^2 + f^2(r)d\vartheta^2\ )\,,$$

$$N(g) = ([0,\infty) \times S^{n-1}\,,\ dr^2 + g^2(r)d\vartheta^2\ )\,,$$

where $f, g \in C^2([0,\infty))$ satisfy the conditions in (1.3). The Euclidean space and the hyperbolic space are corresponding to $f(r) = r$ and $g(r) = \sinh r$, respectively. A map $F : M^n(f) \to M^n(g)$ is called a rotationally symmetric map if

$$F(r,\vartheta) = (\alpha(r)\,,\vartheta) \quad \text{for} \quad \text{all} \quad r > 0 \quad \text{and} \quad \vartheta \in S^{n-1}\,,$$

where $\alpha : [0, \infty) \to [0, \infty)$ with $\alpha(0) = 0$. The energy density [4] of a rotationally symmmetric map $F : M^n(f) \to M^n(g)$ is given by

$$\theta(r) = (\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}\,. \tag{1.2}$$

Given $p > 0$, the p-energy functional [13, 14] for C^2 rotationally symmetric maps is given by

$$\int_0^\infty \left\{ \alpha'^2(r) + (n-1)\frac{g^2(\alpha(r))}{f^2(r)} \right\}^{\frac{p}{2}} f^{n-1}(r)dr\,. \tag{1.3}$$

It can be shown that equation the Euler-Lagrange equation of the p-energy functional is

$$\begin{aligned} \theta^{\frac{p}{2}-1}(r)\alpha''(r) \quad &+ \quad [(n-1)\theta^{\frac{p}{2}-1}(r)\frac{f'(r)}{f(r)} + (\theta^{\frac{p}{2}-1})'(r)]\alpha'(r) \\ &\qquad\qquad -(n-1)\theta^{\frac{p}{2}-1}(r)\frac{g(\alpha(r))g'(\alpha(r))}{f^2(r)} = 0\,. \end{aligned} \tag{1.4}$$

Critical points of the p-energy functional (1.4) are known as rotationally symmetric p-harmonic map from $M(f)$ to $N(g)$. We remark that the notion of p-energy, which generalizes the usual energy as in harmonic maps [4], was first introduced by K. Uhlenbeck [13]. The existence question has been studied later by various authors, using the direct method of Calculus of Variations. In [14], White studied maps minimizing the p-energy in certain homotopy classes. Xu and Yang [15] considered the existence of p-harmonic maps between spheres, generalizing the works of previous authors on harmonic maps between spheres. It is worth mentioning that the direct method yields normally only finite energy maps. For noncompact manifolds, Nakauchi and Takakuwa [10] study concentration compactness and gap theorems for p-harmonic maps.

For $p = 2$, equation (1.4) becomes

$$\alpha''(r) + (n-1)\frac{f'(\alpha)}{f(\alpha)}\alpha'(r) - (n-1)\frac{g(\alpha(r))g'(\alpha(r))}{f^2(\alpha)} = 0\,. \tag{1.5}$$

Equation (1.5) corresponds to rotationally symmetric harmonic maps, which have been studied by Ratto and Rigoli [11] and Cheung and Law [1]. In this note we discuss the existence and properties of positive solutions $\alpha \in C^2((0, \infty)$ to equation (1.4) with

$$\lim_{r \to 0^+} \alpha(r) = 0\,.$$

The results in this note are contained in the works of Cheung-Law-Leung [2] and also in [8, 9].

We first show that any positive solution must be increasing. Equation (1.4) is a quasilinear second order differential equation, where the nonlinear term $\theta^{p/2-1}(r)$ makes the equation complicated. We introduce a first order equation on $\theta(r)$ and $\alpha(r)$ [equation (2.3)]. From this we study unique continuation of nonnegative solutions to equation (1.4). For $p > 2$, we show that if $\alpha(r) \in C^2(0, \epsilon)$ is a nonnegative solution to (1.4) with $\lim_{r\to 0^+} \alpha(r) = 0$, and $\alpha(r)$ goes to zero at arbitrary high order, then $\alpha \equiv 0$ on $(0, \epsilon)$ (lemma 2.6). This unique continuation property is well-known for harmonic maps [7] but is unknown to us for p-harmonic maps in general with $p \neq 2$ (c.f. [6]). By refining the argument, we can estimate the order that α can approach zero. We obtain a Liouville's type theorem for certain rotationally symmetric p-harmonic maps into the Euclidean space or the hyperbolic space (theorem 2.17 & 2.23).

The differential equation (1.4) is solved locally near zero by solving an initial value problem for small r (theorem 3.1). This is achieved by rewriting the differential equation (1.4) as a system of two integral equations and solving it by comparing it to the Euler equation. This local solution is shown to be extendible to the entire positive real axis under certain conditions on f and g. We study asymptotic properties of rotationally symmetric p-harmonic maps from the hyperbolic space to itself, that is, $f(r) = g(r) = \sinh r$. The identity map $\mathrm{Id} : M^n(f) \to M^n(f)$ corresponding to $\alpha(r) \equiv r$ is a rotationally symmetric p-harmonic map. We investigate asymptotic properties of a rotationally symmetric p-harmonic map by seeing whether there is a point "above" or "below" the identity map. More precisely, for $p > 2$, we show that if $F : M^n(\sinh r) \to M^n(\sinh r)$ is a rotationally symmetric p-harmonic map from the hyperbolic space to itself and if there is a point $r_o > (\ln 3)/2$ such that $\alpha(r_o) < r_o$ and $\alpha'(r_o) < 1$, then α is a bounded function. On the other hand if $\alpha(r_o) > r_o$ and $\alpha'(r_o) > 1$, then for all positive number C, $\alpha(r) > r + C$ for all r large. If there is a positive constant C such that $r - C \leq \alpha(r) \leq r + C$ for all r large, then α is asymptotic to the line $y = x$ (theorem 4.18).

We consider the asymptotic properties of positive solutions to (1.4) when $M(f)$ is similar to the hyperbolic space and $N(g)$ is similar to the Euclidean space. we show that $\alpha(r)$ either goes to infinity exponentially or it is bounded (theorem 5.12). If g grows at most exponentially, then we show that either α grows at least linearly or α is bounded (theorem 5.19). We show that, under certain conditions on f and g, all rotationally symmetric p-harmonic maps with $p > 2$ are bounded (corollary 3.24).

2. Unique continuation and Liouville's type theorem

We begin this section with local considerations of equation (1.1).

Lemma 2.1. **[2]** *Let $y(r) \in C^2(0,R)$ be a positive solution to (1.1) with $\lim_{r\to 0^+} y(r) = 0$, where R is a positive number. Then $y'(r) > 0$ for all $r \in (0,R)$.*

Proof. Since $y(r) > 0$ for all $r > 0$ and $\lim_{r\to 0^+} y(r) = 0$, we can find a point r_o such that $y'(r_o) > 0$. Suppose that there exists a point $r_1 > r_o$ such that $y'(r_1) \le 0$, then there exists a point $r' \in (r_o, r_1]$ such that $y(r') > 0$, $y'(r') = 0$ and $y''(r') \le 0$. At r', equation (1.1) gives

$$\theta^{\frac{p}{2}-1}(r')y''(r') = (n-1)\theta^{\frac{p}{2}-1}(r')\frac{g(y(r'))g'(y(r'))}{f^2(r')}\,,$$

and

$$\theta(r') = (n-1)\frac{g^2(y(r'))}{f^2(r')} + (y'(r'))^2 > 0\,.$$

Thus $y''(r') > 0$, contradiction. Therefore $y'(r) > 0$ for all $r \in (r_o\,, R)$. The proof is completed by noticing that we can let $r_o \to 0$. **Q.E.D.**

For $p > 0$, let

$$\Theta(r) = \theta^{\frac{p}{2}-1}(r)\,, \qquad r > 0\,. \tag{2.2}$$

Then

$$\Theta'(r)\theta = (\frac{p}{2}-1)\Theta(r)[(n-1)(\frac{g^2(\alpha(r))}{f^2(r)})' + 2\alpha'(r)\alpha''(r)]\,.$$

Using equation (1.1) to replace the term $\Theta(r)\alpha''(r)$ we have

$$\begin{aligned} &\Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \\ = \;& (p-2)(n-1)\Theta(r)\{\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\}\,. \end{aligned} \tag{2.3}$$

Equation (2.3) involves only first derivatives of Θ and α. For some $\epsilon > 0$, assume that a nonnegative C^2-solution α of (1.4) exists on $(0,\epsilon)$ with $\lim_{r\to 0^+}\alpha(r) = 0$. Given $\kappa > 0$, we can find positive constants a_o, b_o and c_o depending on κ such that

$$|f'(r)| \le a_o\,, \quad f(r) \ge c_o r \quad \text{and} \quad |g'(y(r))| < b_o \quad \text{for} \quad \text{all} \quad r \in (0, \frac{\kappa}{\kappa+1}\epsilon)\,.$$

Then (2.3) gives

$$\begin{aligned} &\Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \\ \le \;& (p-2)(n-1)\Theta(r)\frac{(a_o+b_o)}{c_o r}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\,. \end{aligned} \tag{2.4}$$

Therefore there is a positive constant $C = C(a_o, b_o, c_o, p, n, \kappa)$ such that for $p > 2$ we have

$$\Theta'(r) \leq \frac{C}{r}\Theta(r) \qquad \text{on} \quad (0, \frac{\kappa}{\kappa+1}\epsilon)\,. \tag{2.5}$$

We have the following unique continuation property for solutions of (1.1).

Lemma 2.6. **[9]** *For $p > 2$, let $\alpha(r) \in C^2(0,\epsilon)$ be a nonnegative solution to (1.1) with $\lim_{r\to 0^+}\alpha(r) = 0$. If $\alpha(r) = O(r^k)$ near 0 for all $k > 0$, then $\alpha \equiv 0$ on $(0,\epsilon)$*

Proof. Assume that $\alpha \not\equiv 0$ on $(0, \epsilon/2)$. We first show that α cannot be zero on $(0,\delta)$ for any $\delta \in (0,\epsilon/2)$. Suppose that $\alpha \equiv 0$ on $(0,\delta)$ for some $\delta \in (0,\epsilon/2)$. Since $\alpha \not\equiv 0$ on $(0,\epsilon/2)$, we may assume that $\alpha(r) > 0$ on $(\delta, \epsilon/2)$. Integrating (2.5) we have

$$\ln \Theta(r)|_b^a \leq C \ln r|_b^a\,,$$

where $\delta < b < a < \epsilon/2$. That is

$$\Theta(a) \leq \Theta(b)(\frac{a}{b})^C\,. \tag{2.7}$$

Let $b \to \delta > 0$, we have $\Theta(b) \to 0$, but $\Theta(a) > 0$, contradicting (2.7). Thus α cannot be zero on $(0,\delta)$ for any $\delta \in (0,\epsilon/2)$. Thus we can find a point $r_o \in (0,\epsilon/2)$ such that $\alpha(r_o) > 0$ and $\alpha'(r_o) > 0$. Suppose that there exists a point $r_1 \in (0,\epsilon/2)$ $r_1 > r_o$ such that $\alpha(r_1) = 0$, then there exists a point $r' \in (r_o, r_1)$ such that $\alpha(r') > 0$, $\alpha'(r') = 0$ and $\alpha''(r) \leq 0$. At r', equation (1.4) gives

$$\theta^{\frac{p}{2}-1}(r')\alpha''(r') = (n-1)\theta^{\frac{p}{2}-1}(r')\frac{g(\alpha(r'))g'(\alpha(r'))}{f^2(r)}\,,$$

and

$$\theta(r') = (n-1)\frac{g^2(\alpha(r'))}{f^2(r')} + (\alpha'(r'))^2 > 0\,.$$

Therefore we have $\alpha''(r') > 0$, contradiction. Therefore $\alpha(r) > 0$ for all $r \in (r_o, \epsilon/2)$. As $\lim_{r\to 0^+}\alpha(r) = 0$ and $\alpha \not\equiv 0$ on $(0,\delta)$ for all $\delta > 0$, we can let $r_o \to 0$. Thus $\alpha(r) > 0$ on $(0,\epsilon/2)$. Similarly we can show that $\alpha'(r) > 0$ on $(0,\epsilon/2)$. Integrating (2.5) we obtain

$$\Theta(a) \leq \Theta(r)(\frac{a}{r})^C\,,$$

where in this case $\epsilon/2 > a > r > 0$ and a is a constant. We have

$$\Theta(r) \geq C' r^C \tag{2.8}$$

for some positive constant C' That is,

$$\theta(r) = (n-1)\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2 \geq C'' r^{\frac{2C}{p-2}}\,, \tag{2.9}$$

where C'' is a positive constant. Take a number k such that

$$k > \frac{C}{p-2} + 1\,.$$

Since $\alpha(r) \leq C_k r^k$ for r small, where C_k is a positive constant, and $f(r), g(r) \sim r$ when r is small, we have

$$(n-1)\frac{g^2(\alpha(r))}{f^2(r)} \leq C''' r^{2k-2}\,,$$

where C''' is a positive constant. Hence by (2.8) we have

$$(\alpha'(r))^2 \geq C_o r^{\frac{2C}{p-2}} \tag{2.10}$$

for some positive constant C_o. Therefore we have

$$\alpha(r) - \alpha(b) = \int_b^r \alpha'(s)ds \geq C'' \int_b^r s^{\frac{2C}{p-2}} ds = C_o(r^{\frac{2C}{p-2}+1} - b^{\frac{2C}{p-2}+1})\,, \quad \frac{\epsilon}{2} > r > b > 0\,.$$

As $\lim_{b \to 0^+} \alpha(b) = 0$, we have

$$\alpha(r) \geq C_1 r^{\frac{2C}{p-2}+1}\,, \tag{2.11}$$

contradicting that $\alpha(r) \sim O(r^k)$ for all $k > 0$. Therefore we have $\alpha(r) \equiv 0$ on $(0, \epsilon/2)$. Similar argument shows that $\alpha(r) \equiv 0$ on $0, (\frac{\kappa}{\kappa+1}\epsilon)$ for all $\kappa > 0$. Let $\kappa \to \infty$, we have $\alpha(r) \equiv 0$ on $(0, \epsilon)$. **Q.E.D.**

Let $\alpha \in C^2((0,\epsilon))$ be a nonnegative solution to (1.1) with $\lim_{r \to 0^+} \alpha(r) = 0$. If we assume that $\alpha \not\equiv 0$, then the proof of lemma 2.6 shows that $\alpha(r) > 0$, hence $\alpha'(r) > 0$ on $(0, \epsilon)$ [2]. Given any $\delta > 0$, as $\lim_{r \to 0^+} \alpha(r) = 0$, we can find $\epsilon_o < \epsilon$ such that

$$0 < \alpha(r) < \delta \quad \text{on} \quad (0, \epsilon_o)\,.$$

As $f, g \in C^2(0, \infty)$ with $f'(0) = g'(0) = 1$, given $\varepsilon_1 \in (0,1)$, we can assume that δ and ϵ_o are small enough such that

$$0 < f'(r) \leq 1 + \varepsilon_1\,, \quad f(r) \geq (1-\varepsilon_1)r \ \text{ and } \ |g'(y(r))| < 1 + \varepsilon_1 \quad \text{for} \quad \text{all} \ \ r \in (0, \epsilon_o)\,.$$

If we assume that $p > 2$ and $n \geq 2$, then (2.2) gives

$$\Theta'(r) \leq (p-2)(n-1)(\frac{1+\varepsilon_1}{1-\varepsilon_1})\Theta(r) \quad \text{on} \quad (0, \epsilon_o)\,. \tag{2.12}$$

That is, $C = (p-2)(n-1)(1+\varepsilon_1)/(1-\varepsilon_1)$ in (2.4). It follows as in the proof of lemma 2.5 and inequality (2.9) that

$$\alpha(r) \geq C_1 r^{\frac{2C}{p-2}+1} = C_1 r^{2(n-1)(\frac{1+\varepsilon_1}{1-\varepsilon_1})+1} \,. \tag{2.13}$$

As ε_1 can be made arbitrarily small, therefore we have the following refinement of lemma 2.6.

Theorem 2.14 [9] *For $p > 2$ and $n \geq 2$, let $\alpha(r) \in C^2(0,\epsilon)$ be a nonnegative solution to (1.1) with $\lim_{r\to 0^+} \alpha(r) = 0$. If there exist constants $k > 2n-1, C' > 0$ and $\epsilon_o \in (0,\epsilon)$ such that , $\alpha(r) \leq C' r^k$ for all $r \in (0,\epsilon_o)$, then $\alpha \equiv 0$ on $(0,\epsilon)$*

Lemma 2.15. **[2]** *For $p > 2$ and $n \geq 2$, let $\alpha(r) \in C^2(0,R)$ be a positive solution to (1.1) with $\lim_{r\to 0^+} \alpha(r) = 0$, where R is a positive number. Suppose that f and g satisfy the conditions in (1.3) and $|f'(r)| \leq a$ and $|g'(\alpha(r))| \leq b$ for all $r \in (0,R)$, where $a, b > 0$ are constants, then the relative energy density $\theta(r)$ satisfies the following differential inequality on $(0,R)$:*

$$-\frac{(n-1)(p-2)(a+b)}{(p-1)}\,\frac{1}{f(r)} \leq \frac{(\theta^{q-1})'(r)}{\theta^{q-1}(r)} \leq \frac{(n-1)(p-2)(a+b)}{\min\{p-1\,,\ n-1\}}\,\frac{1}{f(r)}\,. \tag{2.3}$$

Proof. By (2.3) we have

$$\begin{aligned}
&(\theta^{q-1})'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \\
= \ & (p-2)(n-1)\theta^{q-1}(r)\{\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\}\,.
\end{aligned}$$

As $\alpha' > 0$, we have

$$\begin{aligned}
&\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
\leq \ & \frac{1}{f(r)}\{b[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] + a[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\} \\
\leq \ & \frac{1}{f(r)}(a+b)[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\,.
\end{aligned} \tag{2.16}$$

Therefore we have

$$(\theta^{q-1})'(r) \leq C'\frac{\theta^{q-1}(r)}{f(r)}$$

for $r \in (0,R)$, where $C = (p-2)(n-1)(a+b)/\min\{p-1\,,\ n-1\}$ is a positive constant. On the other hand, by (2.4) we have

$$\begin{aligned}
(p-1)\theta(\theta^{q-1})' \ &\geq \ -\frac{(n-1)(p-2)(a+b)}{f(r)}\,\theta^{q-1}[(\alpha')^2 + \frac{g(\alpha)^2}{f(r)^2}]\ , \\
&\geq \ -\frac{(n-1)(p-2)(a+b)}{f(r)}\,\theta^q\ .
\end{aligned}$$

Thus

$$(\theta^{q-1})' \geq -\frac{2(n-1)(q-1)(a+b)}{(p-1)f(r)}\,\theta^{q-1}\ .$$

Q.E.D.

For $p > 2$ and $n \geq 2$, let $F = (\phi, \alpha(r))$ be a rotationally symmetric p-harmonic map from a model space $M^n(f)$ to $N^n(g)$, with $\alpha(r) \geq 0$ defined on $(0, \infty)$ and $\lim_{r\to 0^+} \alpha(r) = 0$, where f and g satisfy the condition in (1.3). We have the following Liouville's type theorem for p-harmonic maps.

Theorem 2.17. **[2]** *Assume that $|f'(r)| \leq c_o^2$, $g'(r) > 0$ and $g'^2(r) - g(r)g''(r) \geq c^2$ for all $r > 0$, where c_o and c are positive numbers. If α is bounded, then $\alpha \equiv 0$.*

Remark. In the Euclidean and hyperbolic case, we have $g(r) = r$ and $g(r) = \sinh r$, respectively. In both cases, we have $g'^2(r) - g(r)g''(r) = 1$ and hence the condition in the above theorem is satisfied. The condition on f is satisfied for the type of functions $f(r) = r^\alpha + \sin r$ for $r \gg 1$ and $\alpha \in (0, 1]$ (c.f. [10] for Liouville's type theorems for harmonic maps).

Proof. Assume that α is bounded and $\alpha \not\equiv 0$, then lemma the proof of lemma 2.6 shows that $\alpha(r) > 0$, $\alpha'(r) > 0$ and hence $\theta(r) > 0$ for all $r > 0$. The p-harmonic map equation is given by

$$(\theta^{q-1})'\alpha' + \theta^{q-1}\alpha'' + (n-1)\frac{f'}{f}\theta^{q-1}\alpha' - (n-1)\frac{\theta^{q-1}}{f^2}g(\alpha)g'(\alpha) = 0\,.$$

Since y is bounded, we may assume that $|g'(\alpha(r))| < a$ for some constant $a > 0$. As in (2.16) we have

$$(\theta^{q-1})'(r) < C\frac{\theta^{q-1}(r)}{f(r)}$$

for $r > 0$, where $C = (p-2)(n-1)(a + c_o^2)/\max\{p-1\,,\ n-1\} + 1$ is a positive constant. We have

$$C\frac{\theta^{q-1}}{f}\alpha' + \theta^{q-1}\alpha'' + (n-1)\frac{f'}{f}\theta^{q-1}\alpha' - (n-1)\frac{\theta^{q-1}}{f^2}g(\alpha)g'(\alpha) > 0\ .$$

Since $\theta \neq 0$ except at the origin, this simplifies to

$$\frac{\alpha''}{g(\alpha)g'(\alpha)} + \frac{n-1}{f}(C + f')\,\frac{\alpha'}{g(\alpha)g'(\alpha)} - (n-1)\frac{1}{f^2} > 0\,, \tag{2.18}$$

for all $r \geq 1$. For $r \geq 1$, let $H = \alpha'/\,[g(\alpha)g'(\alpha)]$. We have

$$\frac{\alpha''}{g(\alpha)g'(\alpha)} = H' + H^2[g'^2(\alpha) + g(\alpha)g''(\alpha)]\,.$$

Since α is a bounded positive solution, we can find a positive constant $C_o > 1$ such that $C_o^2 \geq g'^2(\alpha) + g(\alpha)g''(\alpha)$ for all $r \geq 1$. Thus (2.18) gives

$$C_o^2 H^2 + H' + \frac{n-1}{f}(C + f')\, H - \frac{(n-1)}{f^2} > 0. \tag{2.19}$$

Consider the quadratic form

$$C_o^2 H^2 + \frac{n-1}{f}(C + f')H - \frac{(n-1)}{f^2}\ .$$

It is nonpositive for $H(r) \in [0, b_C(r)]$ where $b_C(r)$ is given by

$$b_C(r) = \frac{n-1}{2fC_o^2}\{[(C+f')^2 + \frac{4C_o^2}{n-1}]^{1/2} - (C + f')\} > 0\,. \tag{2.20}$$

Hence

$$H'(r) > 0 \quad \text{ whenever } \quad H(r) \leq b_C(r)\ . \tag{2.21}$$

Consider the function $q(x) = \sqrt{x^2 + b^2} - x > 0$, where b is a positive number. If $x \leq 0$, then $q(x) \geq b$. For $x > 0$, we have $q'(x) < 0$. Thus $q(x) \geq c_1^2$ if $|x|$ is bounded, where c_1 is a positive number. As $|f'| \leq c_o^2$, we have $|C + f'| \leq C + c_o^2$ and $f(r) \leq C_1 r$ for some positive constant $C_1 > 0$, we can find a positive number C_2 such that

$$b_C(r) \geq \frac{C_2}{r}\,.$$

Thus

$$\int_1^\infty b_C(r)dr = \infty\,.$$

On the other hand we have

$$-(\ln \frac{g'(\alpha)}{g(\alpha)})' = H[g'^2(\alpha) - g(\alpha)g''(y)] \geq c^2 H$$

for $r \geq 1$. Thus

$$-\ln \frac{g'(\alpha(r))}{g(\alpha(r))}|_1^\infty = -\int_1^\infty (\ln \frac{g'(\alpha)}{g(\alpha)})'dr \geq c^2 \int_1^\infty H(r)dr\,.$$

As $g'(r) > 0$ for all $r > 0$ and y is bounded and positive for $r \geq 1$, we have

$$\int_1^\infty H(r)dr < \infty\ .$$

Since $H \geq 0$ and $\int_1^\infty b_C(r)dr = \infty$, for some sufficiently large r we have $H(r) \leq b_C(r)$ and $H'(r) \leq 0$ (c.f. [6]), contradicting (2.21). **Q.E.D.**

It follows from (2.3) that if $f'(r) \geq 0$, then we have

$$\Theta'(r)[(p-1)(\alpha'(r))^2+(n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \leq (p-2)(n-1)\Theta(r)\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)}\,.$$

In addition, if $p > 2$ and and $g'(r) \leq ag(r)$ for all $r > 0$, where a is a positive constant, then there exists a constant C_o such that

$$\Theta'(r) \leq C_o\Theta(r)\alpha'(r)\,. \tag{2.22}$$

We use this to proof the following Liouville's type theorem.

Theorem 2.23 [9] *Assume that there exists positive constants c_o and c_1 such that $0 \leq f'(r) \leq c_o^2$ and $f(r) \leq c_1$ for all $r \geq r_o$. For $p > 2$, let $F(r,\vartheta) = (\alpha(r)\,,\vartheta)$ is a rotationally symmetric p-harmonic maps from $M(f)$ to the Euclidean space or the hyperboic space. If there exists a positive constant C' such that $\alpha'(r) \leq C'$ for all $r > 0$, then $\alpha \equiv 0$.*

Remarks. In fact we prove a more general statement. The conditions below on g are satisfied for the Euclidean and hyperbolic metrics, where $g(r) = r$ and $g(r) = \sinh r$, respectively. Conic type metrics, where $f(r) = k$ for all r large, satisfy the condtions on f in the above theorem.

Proof. If $\alpha \not\equiv 0$, if follows from the proof of lemma 2.6 (c.f. [2]) that $\alpha > 0$ and hence $\alpha'(r) > 0$ for all $r > 0$. Assume that

$$g'(r) \leq ag(r) \quad \text{for} \quad \text{all} \;\; r > 0\,. \tag{I}$$

Then (2.22) implies that

$$\Theta'(r) < C\Theta(r)\alpha'(r)\,,$$

where we may take $C = C_o + 1$. Substitute the above inequality into (1.4) and we have

$$C\Theta[\alpha'(r)]^2 + \Theta\alpha''(r) + (n-1)\frac{f'}{f}\Theta\alpha' - (n-1)\frac{\Theta}{f^2}g(\alpha)g'(\alpha) > 0\,. \tag{2.24}$$

As $0 \leq f'(r) \leq c_o^2$, we have

$$\frac{f'(r)}{f(r)} \leq c_1 \quad \text{for} \quad \text{all} \;\; r \geq r_o\,,$$

where c_1 is a positive constant. Since $\Theta(r) > 0$ and $\alpha'(r) < C'$ for all $r > r_o$, we have

$$\frac{\alpha''}{g(\alpha)g'(\alpha)} + C''\frac{\alpha'}{g(\alpha)g'(\alpha)} - (n-1)\frac{1}{f^2} > 0 \tag{2.25}$$

for all $r \geq r_o$, where C'' is a positive constant. For $r \geq r_o$, let

$$H(r) = \frac{\alpha'(r)}{g(y)g'(y)}\,. \tag{2.26}$$

We have

$$\frac{\alpha''}{g(\alpha)g'(\alpha)} = H' + H^2[g'^2(\alpha)(\alpha) + g(\alpha)g''(\alpha)]\,.$$

Substitute into (2.25) we obtain

$$[g'^2(\alpha) + g(\alpha)g''(\alpha)]H^2 + H' + C'H - \frac{(n-1)}{f^2} > 0. \tag{2.27}$$

As $\alpha' > 0$, we have either $\lim_{r\to\infty}\alpha(r) = \infty$ or α is bounded - in the latter case $\alpha \equiv 0$ by theorem 2.17. Therefore we may assume that $\lim_{r\to\infty}\alpha(r) = \infty$. Assume that

$$g''(r) \geq 0 \tag{II}$$

for all $r \geq r_o$. As $g'(0) = 1$, we have $g'(r) \geq 1$ for all $r > 0$. Thus $\lim_{r\to\infty} g(r) = \infty$. Furthermore, assume that for some positive constant c,

$$g'2(r) - g(r)g''(r) \geq c^2 \quad \text{for} \quad \text{all} \quad r > 0\,. \tag{III}$$

Then we have

$$\begin{aligned} 0 \leq [g'^2(\alpha) + g(\alpha)g''(\alpha)]H^2 \quad &= \quad \frac{\alpha'^2[g'^2(\alpha) + g(\alpha)g''(\alpha)]}{g^2(\alpha)g'^2(\alpha)} \\ &\leq \quad \frac{\alpha'^2}{g^2(\alpha)} + \frac{\alpha'^2 g(\alpha)g''(\alpha)}{g^3(\alpha)g''(\alpha)} \quad \to 0 \quad \text{as} \quad r \to \infty\,, \end{aligned}$$

as by assumption (III) $g'^2(\alpha) > g(\alpha)g''(\alpha)$. Similarly we have

$$\lim_{r\to\infty} H(r) = 0\,.$$

Then (2.27) gives

$$\lim_{r\to\infty} H'(r) > k > 0\,.$$

As $H(r) > 0$, we have

$$\int_{r_o}^{r} H(s)ds = O(r)\,. \tag{2.28}$$

On the other hand by (III) we have

$$-(\ln\frac{g'(y)}{g(y)})' = H[g'^2(y) - g(y)g''(y)] \geq c^2 H$$

for all $r \geq r_o$. Thus

$$(2.29) \qquad -\ln \frac{g'(y(r))}{g(y(r))}\Big|_{r_o}^{r} = -\int_{r_o}^{r} (\ln \frac{g'(y)}{g(y)})' dr \geq c^2 \int_{r_o}^{r} H(s) ds \,.$$

Assume that

$$(\mathrm{IV}) \qquad \frac{g'(s)}{g(s)} \geq a' \quad \text{for} \quad \text{all} \quad s \geq 1 \,,$$

where a' is a positive constant, or,

$$(\mathrm{IV}') \qquad \frac{g'(s)}{g(s)} \geq \frac{C'}{s} \quad \text{for} \quad \text{all} \quad s \geq 1 \,.$$

In the first case we have $a' \leq g'(y)/g(y) \leq a$ for all r large enough. Therefore (2.21) implies that

$$\int_{r_o}^{\infty} H(s) ds < \infty \,.$$

Since $\alpha'(r) < C$, which implies that $\alpha(r) \leq C_1 r$ for all $r \geq r_o$, where C_1 is a positive constant. As $\lim_{r\to\infty} \alpha(r) = \infty$, we may assume that $\alpha(r) \geq 1$ for all r large enough. Therefore in the second case we have

$$a \geq \frac{g'(y)}{g(y)} \geq \frac{C'}{\alpha(r)} \geq \frac{C''}{r}$$

for all r large enough, where C'' is a positive constant. We have

$$\int_{r_o}^{r} H(s) ds \leq C'' \ln r$$

for some constant C''. In both cases, we have a contradiction with (2.28). **Q.E.D.**

For $p = 2$, Liouville's type theorems for harmonic maps have been discussed in [11]. In [12], Liouville's type theorems for p-harmonic maps are obtained under assumptions on the curvature of the manifolds. The conditions in theorems 2.17 & 2.23 involve the first derivative of the solution and growth of $f(r)$, which relax assumptions on the curvature.

3. Existence of bounded positive solutions

In this section we show the following local existence theorem for equation (1.4).

Theorem 3.1. **[2]** *Let f and g be C^2 functions on $[0, \infty)$ which satisfy the conditions in (1.1). Then for any $\alpha > 0$, there is a unique solution $\alpha \in C^2[0, \epsilon)$ to* *equation (1.4) for some* $\epsilon > 0$*, such that* $\alpha(0) = 0$*,* $\alpha'(0) = \alpha$ *and* $\alpha > 0$ *on* $(0, \epsilon)$*.*

Proof. Equation (1.4) is equivalent to

$$\alpha'' + \frac{n-1}{r}\alpha' - \frac{n-1}{r^2}\alpha = ((n-1)(\frac{1}{r} - \frac{f'}{f}) - \frac{(\theta^{q-1})'}{\theta^{q-1}})\,\alpha' + (n-1)(\frac{g(\alpha)g'(\alpha)}{f^2} - \frac{\alpha}{r}) \,.$$

Let $z = \alpha/r$. Then the above equation becomes

$$rz'' + (n+1)z' = [\frac{n-1}{r} - \frac{(n-1)f'}{f} - \frac{(\theta^{q-1})'}{\theta^{q-1}}](rz' + z) + (n-1)(\frac{g(rz)g'(rz)}{f(r)^2} - \frac{z}{r}) \,. \tag{3.2}$$

For equation (3.2), the homogeneous part has linearly independent solutions 1 and r^{-n}, with Wronskian $(-n)r^{-n-1}$. Letting $v(r) = \sqrt{r}z'(r)$, the second order equation (3.2) with $\lim_{r\to 0+} z(r) = \alpha$ is equivalent to the following system of integral equations:

$$\begin{aligned} z(r) &= T_1(z, v) = \alpha + (n-1)\int_0^r (1 - \frac{s^n}{r^n})\Phi(s, v(s), z(s))\,ds \quad \text{and} \\ v(r) &= T_2(z, v) = n(n-1)\int_0^r \frac{s^n}{r^{n+\frac{1}{2}}}\Phi(s, v(s), z(s))\,ds\,, \end{aligned} \tag{3.3}$$

where

$$\Phi(s, v, z) = (\frac{1}{s} - \frac{f'(s)}{f(s)})(\sqrt{s}v + z) - \frac{(\theta^{q-1})'}{(n-1)\theta^{q-1}}(\sqrt{s}v + z) + (\frac{g(sz)g'(sz)}{f(s)^2} - \frac{z}{s}) \,.$$

We apply the method of successive approximation on (3.3). Except for the term

$$A(s, v, z) = \frac{(\theta^{q-1})'}{\theta^{q-1}} \,,$$

$\Phi(s, v, z)$ is only a polynomial in v and z. Thus we need to make sure that all of Φ, $\frac{\partial \Phi}{\partial z}$ and $\frac{\partial \Phi}{\partial v}$ are continuous near $s = 0$ for any $z > 0$ and v. Consider the term $A(s, v, z)$. By (2.3), we have

$$\begin{aligned} A(s, v, z) &= \frac{2(q-1)(n-1)}{(2q-1)(\sqrt{s}v + z)^2 + (n-1)g(sz)^2/f(s)^2} \left[\frac{2g(sz)g'(sz)(\sqrt{s}v + z)}{f(s)^2} \right. \\ &\qquad \left. - \frac{f'(s)(\sqrt{s}v + z)^2}{f(s)} - \frac{g(sz)^2 f'(s)}{f(s)^3}\right] \,, \end{aligned}$$

where the denominator is a differentiable function and never vanishes when $s > 0$. Assume that when r is close to 0,

$$\begin{aligned} f(r) &= r + f_1 r^2 + o(r^2) \;; \\ g(r) &= r + g_1 r^2 + o(r^2) \;, \end{aligned}$$

for some $f_1, g_1 \in \mathbb{R}$. A computation gives

$$\lim_{s\to 0} A(s,v,z) = \frac{2(q-1)(n-1)}{(2q+n-2)z^2}\left(-v^2 + 4g_1 z^3 - 4f_1 z^2\right), \tag{3.4}$$

and

$$\begin{aligned} \lim_{s\to 0}\frac{\partial A}{\partial v}(s,v,z) &= \frac{-4(q-1)(n-1)v}{(2q+n-2)z^2} = \frac{\partial A}{\partial v}(0,v,z)\,, \\ \lim_{s\to 0}\frac{\partial A}{\partial z}(s,v,z) &= \frac{4(q-1)(n-1)(v^2+2g_1 z^3)}{(2q+n-2)z^3} = \frac{\partial A}{\partial z}(0,v,z)\,. \end{aligned}$$

It follows that Φ, $\frac{\partial \Phi}{\partial z}$ and $\frac{\partial \Phi}{\partial v}$ are all continuous near $(0,z,v)$ for any $z>0$ and v. Moreover, as s tends to 0,

$$\frac{1}{s} - \frac{f'(s)}{f(s)} = f_1 + O(s)\,, \tag{3.5}$$

and

$$\frac{g(sz)g'(sz)}{f(s)^2} - \frac{z}{s} = 3g_1 z^2 - 2f_1 z + o(1)\,. \tag{3.6}$$

For $\epsilon > 0$ sufficiently small, we consider the complete metric space

$$S = \{(z,v):\ z, v \in C(0,\epsilon) \text{ with } \sup_{(0,\epsilon)} |z(r)-\alpha| \le \frac{\alpha}{2} \text{ and } \sup_{(0,\epsilon)} |v(r)| \le 1\}\,.$$

By equations (3.4), (3.5) and (3.6), for any $(z,v)\in S$ and $\epsilon < \alpha^2/4$,

$$\begin{aligned} & |\Phi(s, v(s), z(s))| \\ \le\ & |f_1|\alpha + \frac{16(q-1)(n-1)}{\alpha(2q+n-2)}(1 + \frac{27}{2}|g_1|\alpha^3 + 9|f_1|\alpha^2) + (\frac{27}{2}|g_1|\alpha + |f_1|)\alpha^2 + \eta\,, \end{aligned}$$

where $\eta > 0$ is small. Thus the operator defined by $T = (T_1, T_2)$ as given in (3.3) maps S into itself. Therefore by contraction mapping theorem, the system (3.3) has a unique solution $(z,v)\in C[0,\epsilon)$ for some $\epsilon > 0$, with $z(0)=\alpha$ and $v(0)=0$. Since $z' = v/\sqrt{r}$, $y' = \sqrt{r}v + z$, and

$$v'(r) = \frac{(n-1)\Phi}{r} - (n+\frac{1}{2})\frac{v}{r}\,, \tag{3.7}$$

we have $z, v\in C^\infty(0,\epsilon)$ and so $y \in C^\infty(0,\epsilon)\cap C^1[0,\epsilon)$. It remains to show that y is C^2 at $r=0$. To do this, we observe that by L'Hopital's rule,

$$\begin{aligned} \lim_{r\to 0} z'(r) &= \lim_{r\to 0}\frac{v}{\sqrt{r}} = \lim_{r\to 0}(n-1)\int_0^r \frac{s^n}{r^{n+1}}\Phi(s,v(s),z(s))\,ds \\ &= \frac{n-1}{n+1}\Phi_0\,, \end{aligned}$$

which is also equal to $z'(0)$. Here

$$\Phi_0 \overset{\text{def}}{=} \lim_{s\to 0}\Phi(s,v(s),z(s)) = \frac{(n-1)(n-2q+2)g_1\alpha^2+(4nq-2n^2-2q-n+2)f_1\alpha}{2q+n-2} .$$

As a result,

$$\alpha''(0) = \lim_{r\to 0}\frac{\alpha'(r)-\alpha}{r} = \frac{2(n-1)}{n+1}\Phi_0 = \lim_{r\to 0}\alpha''(r) .$$

Q.E.D.

Theorem 3.8. **[2]** *For $p>2$, let $\alpha(r)\in C^2(0,R)$ be a positive solution to (1.1) with $\lim_{r\to 0^+}\alpha(r)=0$, where R is a positive number. Suppose that f and g satisfy the conditions in (1.4) and there exists a positive constant a such that $|g'(r)|\le a$ for all $r\ge 0$. Then y can be extended to a positive solution of (1.4) on $(0,\infty)$.*

Proof. By the continuation of solutions [3, p.15], it suffices to show that a solution of equation (1.4) is bounded on the interval $(0,R)$. There exists a positive constant b such that $|f'(r)|\le b$ and $|g'(\alpha(r))|\le a$ for all $r\in(0,R)$. By lemma 2.15 we have $\theta(r)<C$ for all $r\in(R/2,R)$. Thus $\alpha'(r)$ and hence $\alpha(r)$ are bounded on $(R/2,R)$. We can continue the solution to $(0,R+\epsilon)$ for some positive number ϵ. **Q.E.D.**

Theorem 3.9. **[2]** *Suppose that $p>2$ and $n\ge 2$, and assume that there exist constants $a>0$, and $\delta>1$ with $(n-1)\delta>2q-1$ ($q=p/2$) such that*

$$g(y)>0, \qquad 0\le g'(y)\le a \quad \text{for any } y>0 ,$$

and

$$f(r)\asymp r^\delta \qquad f'(r)>0 \quad \text{for large } r ,$$

then all positive solutions to equation (1.1) are bounded if and only if $n>1$.

Remark. When the above conditions are satisfied, by using theorem 3.1 and theorem 3.8, there exists a family of global solutions which are all bounded harmonic maps.

Proof. When $n=1$, then $\theta=(\alpha')^2$. Equation (1.4) implies that α' is constant. Hence

$$\alpha'(r)=\alpha'(0)=\alpha , \qquad \text{or} \qquad \alpha(r)=\alpha r ,$$

which is unbounded. Asume that $n>1$. Equation (1.4) implies that

$$\begin{aligned}(\theta^{q-1}\alpha')' &= (n-1)\theta^{q-1}\Big(\frac{g(\alpha)g'(\alpha)}{f^2}-\frac{f'\alpha'}{f}\Big)\\ &\le (n-1)\theta^{q-1}\Big(\frac{C\alpha}{r^{2\delta}}\Big) ,\end{aligned}$$

when r is sufficiently large, say, when $r \geq r_0$. Hence

$$(\theta^{q-1}\alpha') \leq C\,(\int_{r_0}^{r} s^{-2\delta}\theta^{q-1}\alpha\,ds + 1)\,.$$

Apply Lemma 2.15 to obtain

$$\begin{aligned}(\theta^{q-1}\alpha)' &\leq 2(n-1)(q-1)(a+1)\frac{\theta^{q-1}\alpha}{r^{\delta}} + C\,(\int_{r_0}^{r} s^{-2\delta}\theta^{q-1}\alpha\,ds + 1)\ , \\ &= \frac{2\delta}{r}\,\theta^{q-1}\alpha + C(\int_{r_0}^{r} s^{-2\delta}\theta^{q-1}\alpha\,ds + 1)\end{aligned}$$

for $r \geq r_0$, where r_o is a positive constant. Thus

$$(r^{-2\delta}\theta^{q-1}\alpha)' = Cr^{-2\delta}\left(\int_{r_0}^{r} s^{-2\delta}\theta^{q-1}\alpha\,ds + 1\right)\,. \tag{3.10}$$

Let

$$u = \int_{r_0}^{r} s^{-2\delta}\theta^{q-1}\alpha\,ds + 1\,.$$

Equation (3.10) can be written as

$$u'' \leq Cr^{-2\delta}u \leq \epsilon^2 r^{-2}u \tag{3.11}$$

for some small $0 < \epsilon < 2\delta - 2$. So u is a supersolution of the Euler equation. By the weak maximum principle, we obtain

$$u \leq Cr^{1+\epsilon}\ .$$

Substitute this result into equation (3.10) and integrate. We have

$$r^{-2\delta}\theta^{q-1}\alpha \leq C\,(r^{-2\delta+2+\epsilon} + 1) \leq C$$

for $r \geq r_0$, and so

$$\int_{r_0}^{r} s^{(n-3)\delta}\theta^{q-1}\alpha\,ds \leq Cr^{(n-1)\delta+1}\ . \tag{3.12}$$

On the other hand, we have

$$(f^{n-1}\theta^{q-1}\alpha')' \leq (n-1)af^{n-3}\theta^{q-1}\alpha\,. \tag{3.13}$$

Integrate from r_0 to r, we obtain

$$f^{n-1}\theta^{q-1}\alpha' \leq C\,(\int_{r_0}^{r} s^{(n-3)\delta}\theta^{q-1}\alpha\,ds + 1)\,. \tag{3.14}$$

Substitute equation (3.13) into equation (3.14) to obtain

$$(\alpha')^{2q-1} \leq \theta^{q-1}\alpha' \leq Cr\,.$$

After integration we have

$$\alpha \leq C r^{1+1/(2q-1)}\,,$$

This implies that

$$\theta^{q-1}\alpha \leq C r^{2}\,.$$

Substitute this result into equation (3.14) again,

$$(\alpha')^{2q-1} \leq C r^{3-2\delta} + C r^{-(n-1)\delta}\;.$$

If $2\delta - 2 \geq (n-1)\delta > 2q-1$, then

$$\alpha' \leq C r^{\frac{-(n-1)\delta}{2q-1}}\,.$$

Therefore we have

$$\alpha \leq C\,(r^{1-\frac{(n-1)\delta}{2q-1}} + 1)\,, \tag{3.15}$$

which means that α is bounded, as required. Otherwise,

$$\alpha \leq C\,(r^{1+\frac{3-2\delta}{2q-1}} + 1)\,.$$

When $1 + \frac{3-2\delta}{2q-1} < 0$, then α is bounded. If not,

$$\theta^{q-1}\alpha \leq C r^{4-2\delta}\;.$$

Substitute into equation (3.14),

$$(\alpha')^{2q-1} \leq C r^{5-4\delta} + C r^{-(n-1)\delta}\;.$$

In this way, we have a sequence of upper bounds for $(y')^{2q-1}$ as a linear combination of $r^{2k+1-2k\delta}$ and $r^{-(n-1)\delta}$. Since $\delta > 0$, $r^{-(n-1)\delta}$ will dominate after a finite number of iterations. Therefore equation (3.15) holds eventually. Thus y is bounded. **Q.E.D.**

Theorem 3.16. [**2**] *For $p > 2$ and $n \geq 2$, if $(n-1)\delta \leq 2q-1$, where $q = p/2$, and $f \asymp r^{\delta}$ for large r, then no positive entire solution to equation (1.4) is bounded.*

Proof. Observe by equation (3.13), the function $f^{n-1}\theta^{q-1}y'$ is increasing to some $P > 0$ (may be ∞). Hence there is some constant C such that for large r,

$$\theta^{q-1}\alpha' \geq C r^{-(n-1)\delta}\;. \tag{3.17}$$

Suppose for contradiction that a solution y is bounded. Then we must have $\theta \asymp (\alpha')^{2}$. For if otherwise, we have a sequence of points where

$$\theta \asymp r^{-2\delta} \quad \text{and} \quad \alpha' = o(r^{-\delta})\;.$$

So

$$\begin{aligned} f^{n-1}\theta^{q-1}\alpha' &= o(r^{\delta(n-2q)})\ , \\ &= o(1)\ , \end{aligned}$$

which is impossible, since $P > 0$. Therefore by equation (3.17) we have

$$\alpha' \geq Cr^{\frac{-(n-1)\delta}{2q-1}}$$

for large r, so that y is greater than $Cr^{1-\frac{(n-1)\delta}{2q-1}}$, or $C\log r$. In both cases, y is unbounded. This gives a contradiction.**Q.E.D.**

Theorem 3.18. [**2**] *For $p > 2$ and $n \geq 2$, if $(n-1)\delta \leq 2q-1$, $f \sim Cr^{\delta}$, and $0 < g' \leq a$ $(a > 0)$, then any positive entire solution y of equation (1.4) satisfies*

$$y' \sim Ar^{\frac{-(n-1)\delta}{2q-1}} \qquad \text{for some constant } A > 0\ .$$

Proof. As $f^{n-1}\theta^{q-1}y'$ tends to a number P, finite or infinite. Suppose that $(n-1)\delta < 2q-1$. If $P = \infty$, then we can find a sequence $R_N \to \infty$ such that, at $r = R_N$, $f^{n-1}\theta^{q-1}y'(R_N) = N$, and for $r \leq R_N$,

$$f^{n-1}\theta^{q-1}y'(r) \leq N\ .$$

Hence $r^{(n-1)\delta}(y')^{2q-1} \leq CN$ for some constant C. This gives, after an integration,

$$\alpha(r) \leq CN^{\frac{1}{2q-1}}r^{1-\frac{(n-1)\delta}{2q-1}}\ .$$

Thus

$$\theta(r) \leq CN^{\frac{2}{2q-1}}r^{-\frac{2(n-1)}{2q-1}}\ .$$

So by equation (3.10) we have

$$\begin{aligned} N &= O\left(\int_r^{R_N} \theta^{q-1}N^{\frac{1}{2q-1}}s^{(n-3)\delta+1-\frac{(n-1)\delta}{2q-1}}\,ds + 1\right)\ , \\ &= O\left(N\int_r^{R_N} s^{1-2\delta}\,ds + 1\right)\ , \\ &= o(1)N \end{aligned}$$

for r sufficiently large. Therefore, as $N \to \infty$, $N = o(N)$ which is impossible. Hence there exists some constant $P \in \mathbf{R}$ such that

$$f^{n-1}\theta^{q-1}y' \sim P\ . \tag{3.19}$$

Thus

$$\alpha' \leq Cr^{\frac{-(n-1)\delta}{2q-1}} \quad \text{and} \quad \alpha \leq Cr^{1-\frac{(n-1)\delta}{2q-1}}\ .$$

Substituting these bounds into θ in equation (3.17), we obtain

$$y' \asymp r^{\frac{-(n-1)\delta}{2q-1}} \quad \text{and} \quad \alpha \asymp r^{1-\frac{(n-1)\delta}{2q-1}}$$

Therefore $\theta \sim (\alpha')^2$ and

$$\alpha' \sim Ar^{\frac{-(n-1)\delta}{2q-1}} \qquad \text{for some } A > 0 \ .$$

When $(n-1)\delta = 2q-1$, the procedure is similar.

$$\alpha' \le CN^{\frac{1}{2q-1}} r^{-1} \ ,$$

and

$$\alpha \le CN^{\frac{1}{2q-1}} \log r \ .$$

So

$$N \le CN \int_r^{R_N} s^{1-2\delta} \log s \, ds = o(1)N \ .$$

The other cases follow in the same way. **Q.E.D.**

Corollary 3.20. [2] *For $p > 2$ and $n \ge 2$, if $\delta > \frac{2q-1}{n-1} \ge 1$, $f \sim Cr^\delta$, and $0 < g' \le a$, then*

$$y' \sim Ar^{\frac{-(n-1)\delta}{2q-1}} \qquad \text{for some } A > 0 \ .$$

Proof. We follow the proof in Theorem 3.11. Suppose there is a sequence $R_N \to \infty$ such that $f^{n-1}\theta^{q-1}y'(R_N) = N$, then for $r < R_N$,

$$\alpha' \le CN^{\frac{1}{2q-1}} r^{-\frac{(n-1)\delta}{2q-1}} \ ,$$

but

$$\alpha \le CN^{\frac{1}{2q-1}} \ .$$

Hence

$$\begin{aligned} N &\le O\left(N \int_r^{R_N} s^{(n-3)\delta} \max\{s^{-(n-1)\delta}, \ s^{-2\delta(q-1)}\} \, ds + 1 \right) \\ &= O(N \int_r^{R_N} s^{-\delta} \, ds + 1) \ , \\ &= o(1)N \end{aligned}$$

for r sufficiently large, and as $N \to \infty$. The other cases are similar. **Q.E.D.**

Corollary 3.21. [2] *For $p > 2$ and $n \ge 2$, if $\delta > \frac{2q-1}{n-1} \ge 1$, $f \sim Cr^\delta$, and $0 < g' \le a$, then*

$$\alpha' \sim Ar^{\frac{-(n-1)\delta}{2q-1}} \qquad \text{for some } A > 0 \ .$$

Proof. We follow the proof in Theorem 3.18. Suppose there is a sequence $R_N \to \infty$ such that $f^{n-1}\theta^{q-1}y'(R_N) = N$, then for $r < R_N$,

$$\alpha' \le C N^{\frac{1}{2q-1}} r^{-\frac{(n-1)\delta}{2q-1}} \ ,$$

but

$$\alpha \le C N^{\frac{1}{2q-1}} \ .$$

Hence

$$\begin{aligned} N &\le O\left(N \int_r^{R_N} s^{(n-3)\delta} \max\{s^{-(n-1)\delta},\ s^{-2\delta(q-1)}\}\, ds + 1\right) \\ &= O(N \int_r^{R_N} s^{-\delta}\, ds + 1) \ , \\ &= o(1)N \end{aligned}$$

for r sufficiently large, and as $N \to \infty$. The other cases are similar. **Q.E.D.**

Theorem 3.22. **[9]** *Assume that* $f'(r) > 0$ *for all* $r > 0$ *and* $\lim_{r\to\infty} f'(r) = \infty$. *Suppose that* $g'(y) \le k$ *for all* $y \ge 0$, *where* k *is a positive constant. Let* $\alpha(r) \in C^2(0,\infty)$ *be a positive solution to (1.4) with* f *and* g *as above and* $\lim_{r\to 0^+} \alpha(r) = 0$. *If* $2 < p \le n$, *then for any* $\epsilon \in (0,1)$, *there exist positive constants* r_o *and* C *such that* $\alpha'(r) \le C/f^{1-\epsilon}(r)$ *for all* $r \ge r_o$. *If* $p > n \ge 2$, *then for any* $\epsilon > 0$, *there exist positive constants* r_o *and* C *such that*

$$\alpha'(r) \le \frac{C}{(f(r))^{(1-\epsilon)\frac{(n-1)}{(p-1)}}}$$

for all $r \ge r_o$.

Proof. Given $1 > \epsilon > 0$, as in (5.15) (with $\epsilon = 1 - \tau$) we can find a positive number r_o such that

$$\begin{aligned} &\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\ \le\ & \frac{g^2(\alpha(r))}{\epsilon f^3(r) f'(r)}[k^2 - \epsilon^2 f'(r)^2] - (1-\epsilon)[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \\ \le\ & -(1-\epsilon)\frac{f'(r)}{f(r)}[\frac{\alpha^2(r)}{f^2(r)} + (\alpha'(r))^2] \end{aligned}$$

for all $r \ge r_o$, as $g'(s) \le k$ and $\lim_{r\to\infty} f'(r) = \infty$. By (2.15) we have

$$\begin{aligned} &\Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \\ \le\ & -(1-\epsilon)(p-2)(n-1)\frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\Theta(r) \end{aligned} \tag{3.23}$$

for all $r \geq r_o$. Thus if $2 < p \leq n$, then (3.23) implies that

$$\Theta'(r) \leq -(1-\epsilon)(p-2)\frac{f'(r)}{f(r)}\Theta(r)$$

for all $r \geq r_o$. An integration gives $\alpha'(r) \leq C/f^{1-\epsilon}(r)$ for some positive constant C and for all $r \geq r_o$. If $p > n > 2$, then (3.19) gives

$$\Theta'(r) \leq -(1-\epsilon)\frac{(p-2)(n-1)}{p-1}\frac{f'(r)}{f(r)}\Theta(r)$$

for all $r \geq r_o$. An integration gives

$$\alpha'(r) \leq \frac{C}{(f(r))^{(1-\epsilon)\frac{(n-1)}{(p-1)}}}$$

for some positive constant C and for all $r \geq r_o$. **Q.E.D.**

As a corollary, we have the following result (c.f. [2]).

Corollary 3.24. **[9]** *Assume that $f'(r) > 0$ for all $r > 0$ and $\lim_{r\to\infty} f'(r) = \infty$. Suppose that $g'(y) \leq k$ for all $y > 0$, where k is a positive constant. Assume that $f(r) \geq Cr^s$ for some positive constant C and for all $r > \bar{r} > 0$, where $s > 1$ if $2 < p \leq n$, and*

$$s > \frac{(p-1)}{(n-1)}$$

if $p > n \geq 2$. Let $\alpha(r) \in C^2(0,\infty)$ be a positive solution to (1.1) with f and g as above and $\lim_{r\to 0^+} \alpha(r) = 0$. Then α is a bounded function on $I\!R^+$.

Proof. From theorem 3.21, we can choose $r_o > 0$ and $\epsilon > 0$ so that

$$\alpha'(r) \leq \frac{C}{r^{1+\delta}(r)}$$

for all $r \geq r_o$. Here δ is a positive constant. An integration shows that α is a bounded function on $\mathbb{R}^+$. **Q.E.D.**

In particular, for $p > 2$, all rotationally symmetric p-harmonic maps from the hyperbolic space to the Euclidean space are bounded.

4. p-harmonic maps from hyperbolic space to itself

In this section we discuss rotationally symmetric p-harmonic maps from the hyperbolic space to itself. In this case $f(r) = g(r) = \sinh r$ as the metric

$$dr^2 + \sinh^2 r\, d\varphi^2$$

is exactly the hyperbolic metric. The results in this section can be found in [8].

Lemma 4.1. *For $p > 2$, let $F(r, \varphi) = (\alpha(r), \varphi)$ be a rotationally symmetric p-harmonic map from the hyperbolic space to itself, where $\alpha \in C^2(0, \infty)$ and $\lim_{r\to 0^+} \alpha(r) = \alpha(0) = 0$. If there exists $r_o > (\ln 3)/2$ such that $\alpha(r_o) = r_o + \delta$ for some $\delta \geq 0$ and $\alpha'(r_o) > 1$, then $\alpha(r) > r + \delta$ and $\alpha'(r) > 1$ for all $r > r_o$. If there exists $r_o > (\ln 3)/2$ such that $\alpha(r_o) = r_o - \delta$ for some $\delta \geq 0$ and $\alpha'(r_o) < 1$, then $\alpha(r) < r_o - \delta$ and $\alpha'(r) < 1$ for all $r > r_o$.*

Proof. Suppose that there exists $r_o > (\ln 3)/2$ such that $\alpha(r_o) = r_o + \delta$ for some $\delta \geq 0$ and $\alpha'(r_o) > 1$. Assume that there is a point $r' > r_o$ such that $\alpha(r') = r' + \delta$. Then there exists a point $r \in (r_o, r')$ such that $\alpha(r) > r + \delta$, $\alpha'(r) = 1$ and $\alpha''(r) \leq 0$. Let $\alpha(r) = r + \epsilon$ where $\epsilon > \delta$ is a constant. As $f(r) = g(r) = \sinh r$, at the point r where $\alpha'(r) = 1$ we have

$$\begin{aligned}
& \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
= \;& \frac{1}{8f^3(r)}[2(e^r - e^{-r})(e^\alpha - e^{-\alpha})(e^\alpha + e^{-\alpha}) \\
& \qquad -(e^r + e^{-r})(e^\alpha - e^{-\alpha})^2 - (e^r + e^{-r})(e^r - e^{-r})^2] \\
= \;& \frac{1}{8f^3(r)}(e^r e^{2\alpha} - 3e^r e^{-2\alpha} - 3e^{-r}e^{2\alpha} + e^{-r}e^{-2\alpha} + 3e^r + 3e^{-r} - e^{-3r} - e^{3r})\,.
\end{aligned}$$

As $\alpha(r) = r + \epsilon$, we have

$$\begin{aligned}
(4.2) \qquad & \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
= \;& \frac{1}{8f^3}[(e^{2\epsilon} - 1)(e^{3r} - 3e^r) + (3e^{-r} - e^{-3r})(1 - e^{-2\epsilon})]\,.
\end{aligned}$$

If $r > (\ln 3)/2$, then $e^{3r} - 3e^r > 0$ and $3e^{-r} - e^{-3r} > 0$. Equation (4.2) implies that

$$\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] > 0$$

and by (2.3) $\Theta'(r) > 0$ because $\theta(r) > 0$. As

$$\Theta'(r) = (\frac{p}{2} - 1)\theta^{\frac{p}{2}-2}(r)\theta'(r)\,,$$

hence $\theta'(r) > 0$ if $p > 2$. We have

$$\theta'(r) = (n-1)(\frac{g^2(\alpha)}{f^2(r)})' + 2\alpha'(r)\alpha''(r)\,.$$

And

$$(4.3) \qquad \begin{aligned} (\frac{g^2(\alpha)}{f^2(r)})' &= \frac{2f(r)g(\alpha)[f(r)g'(\alpha)\alpha'(r) - f'(r)g(\alpha)]}{f^4(r)} \\ &= \frac{g(\alpha)}{f^3(r)}(e^r e^{-\alpha} - e^{-r}e^{\alpha})\,. \end{aligned}$$

As $\alpha(r) > r$, we have

$$(\frac{g^2(\alpha)}{f^2(r)})' < 0\,.$$

Hence $\alpha''(r) > 0$. This is a contradiction. Hence $\alpha(r) > r + \delta$ for all $r > r_o$. Suppose that there is a point r' such that $\alpha(r') = 1$, then there is a point $r \in (r_o, r']$ such that $\alpha(r) > r + \delta$, $\alpha'(r) = 1$ and $\alpha''(r) \le 0$. This would give us the same contradiction. Hence $\alpha'(r) > 1$ for all $r > r_o$. Similar argument works for the case $\alpha(r_o) = r_o - \delta$. **Q.E.D.**

Lemma 4.4. *For $p > 2$, let $F(r, \varphi) = (\alpha(r), \varphi)$ be a rotationally symmetric p-harmonic map from the hyperbolic space to itself, where $\alpha \in C^2(0, \infty)$ and $\lim_{r\to 0^+} \alpha(r) = \alpha(0) = 0$. Then we can find a point $r_o > 0$ such that either $\alpha(r) \ge r$ or $\alpha(r) \le r$ for all $r > r_o$.*

Proof. Suppose that we cannot find such a point r_o. Then there exist two points $r_2 > r_1 > (\ln 3)/2$ such that $\alpha(r_1) = r_1$ and $\alpha(r_2) = r_2$ and $\alpha(r) < r$ or $\alpha(r) > r$ for all $r \in (r_1\,, r_2)$. Then we can find a point $r \in (r_1\,, r_2)$ such that $\alpha'(r) = 1$ and if $\alpha(r) > r$, then $\alpha''(r) \le 0$; if $\alpha(r) < r$, then $\alpha''(r) \ge 0$. But the proof of lemma 4.1 shows that these are impossible. **Q.E.D.**

We have the following Liouville's type theorem for p-harmonic map on hyperbolic spaces.

Theorem 4.5. *For $p > 2$, let $F(r, \varphi) = (\alpha(r), \varphi)$ be a rotationally symmetric p-harmonic map from the hyperbolic space to itself, where $\alpha \in C^2(0, \infty)$ and $\lim_{r\to 0^+} \alpha(r) = \alpha(0) = 0$. Assume that there exists $r_o > (\ln 3)/2$ such that $\alpha(r_o) >$*

r_o and $\alpha'(r_o) > 1$. Then given any positive number C, we have a point r_c such that $\alpha(r) > r + C$ for all $r > r_c$.

Proof. Assume that there is a positive number C such that $\alpha(r) \le r + C$ for all $r > r_o$. Let $\alpha(r_o) = r_o + \epsilon_o$ for some constant $\epsilon_o > 0$. By lemma 4.1, we have $\alpha(r) > r + \epsilon_o$ and $\alpha'(r) > 1$ for all $r \ge r_o$. Assume that at a point $r > \bar{r}$, $\alpha(r) = r + \epsilon$ and $\alpha'(r) \le 1 + \eta$, where $\epsilon \ge \epsilon_o$, and $\bar{r} > r_o$ and η are a positive constants to be determined later. Then

$$\begin{aligned}
& \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
\ge\ & \frac{1}{8f^3(r)}[2(e^{3r}e^{2\epsilon} - e^{-r}e^{-2\epsilon} - e^{r}e^{2\epsilon} + e^{-3r}e^{-2\epsilon}) \\
& -(e^{3r}e^{2\epsilon} - 2e^{r} + e^{-r}e^{-2\epsilon} + e^{r}e^{2\epsilon} - 2e^{-r} + e^{-3r}e^{-2\epsilon}) \\
& -(1+\eta)^2(e^{3r} - e^{r} - e^{-r} + e^{-3r})] \\
=\ & \frac{1}{(e^r - e^{-r})^3}\{e^{3r}[e^{2\epsilon} - (1+\eta)^2] \\
& -e^{r}[3e^{2\epsilon} - 2 + (1+\eta)^2] - e^{-r}[3e^{2\epsilon} - 2 + (1+\eta)^2] \\
& -e^{-3r}[e^{-2\epsilon} - (1+\eta)^2]\}\,.
\end{aligned}$$

Therefore if we choose $\bar{r} > r_o$ to be large and fix η such that

$$e^{2\epsilon_o} > (1+\eta)^2 + \frac{(e^{2\epsilon_o} - 1)}{2}\,, \tag{4.6}$$

then there exists a positive constant $c > 0$ such that

$$\frac{2g(\alpha)g'(\alpha)\alpha'}{f^2} - \frac{f'}{f}[\frac{g^2(\alpha)}{f^2(r)} + (\alpha')^2] \ge c$$

whenever $r > \bar{r}$ and $\alpha'(r) \le 1 + \eta$. That is,

$$\Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha)}{f^2(r)}] \ge c\Theta(r)$$

for all $r > \bar{r}$ with $\alpha'(r) \le 1 + \eta$. As $\alpha'(r) \le 1 + \eta$ and $\alpha(r) \le r + C$ for some positive constant C, we have

$$\begin{aligned}
& (p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha)}{f^2(r)} \\
\le\ & (p-1)(1+\eta)^2 + \frac{(n-1)}{4}\frac{e^{2C} - 2e^{-2r} + e^{-4r}e^{-2C}}{1 - 2e^{-2r} + e^{-4r}} \le c'\,,
\end{aligned} \tag{4.7}$$

where c' is a positive constant. Hence

$$\Theta'(r) \ge c_o\Theta(r) \quad \text{for all} \quad r > \bar{r} \quad \text{with} \quad \alpha'(r) \le 1 + \eta\,, \tag{4.8}$$

where $c_o = c/c'$ is a positive constant. Since the set $O = \{r > \bar{r} |\ \alpha'(r) > 1 + \eta\}$ is an open set, we can decompose O as

$$O = \cup_{i=1}^{\beta} (a_i, b_i)\,,$$

where $\beta \in \mathbf{N} \cup \{\infty\}$ and $a_i < b_i < a_{i+1}$. For $r \in (a_i, b_i)$, we have $\alpha'(r) > 1 + \eta$ and

$$(\frac{g^2(\alpha)}{f^2(r)})' \geq \frac{f(r)g^2(\alpha)}{2f^4(r)}[\eta e^r e^{\alpha} + (2+\eta)e^r e^{-\alpha} - (2+\eta)e^{-r}e^{\alpha} - \eta e^{-r}e^{-\alpha}] \geq 0$$

if we choose $\bar{r}$ to be large. Since $\alpha'(a_i) = \alpha'(b_i) = 1 + \eta$ and on (a_i, b_i) we have

$$(\frac{g^2(\alpha)}{f^2(r)})' \geq 0\,,$$

therefore

$$\theta(a_i) = (n-1)\frac{g^2(\alpha(a_i))}{f^2(a_i)} + [\alpha'(a_i)]^2 \leq (n-1)\frac{g^2(\alpha(b_i))}{f^2(b_i)} + [\alpha'(b_i)]^2 = \theta(b_i)\,.$$

That is,

$$\Theta(a_i) \leq \Theta(b_i) \tag{4.9}$$

for all $i = 1, 2, ..., \beta$. On the open set O we have $\alpha' > 1 + \eta$. As $\alpha(r) > 1$ for all $r > r_o$, by lemma 4.1, we have

$$r + C \geq \alpha(r) \geq r + \eta[\sum_{i=1}^{\beta(r)} (b_i - a_i)]\,,$$

where $\beta(r)$ is the biggest index j such that $b_j < r$. Therefore

$$C \geq \eta \sum_{i}^{\beta} (b_i - a_i)\,.$$

So the compliment of O on (a_1, ∞), which is given by $U_{i=1}^{\beta}[b_i, a_{i+1}]$, has infinite measure, where $a_{\beta+1} = \infty$ if β is an finite integer. For $r \in [b_i, a_{i+1}]$ we have $\alpha'(r) \leq 1 + \eta$ and by (4.5), $\Theta'(r) \geq c_o\Theta(r)$, or

$$\Theta(a_{i+1}) \geq \Theta(b_i)e^{c_o(a_{i+1}-b_i)}$$

for all i. Apply (4.6) we obtain

$$\begin{aligned}
\Theta(a_{i+1}) &\geq \Theta(b_i)e^{c_o(a_{i+1}-b_i)} \\
&\geq \Theta(a_i)e^{c_o(a_{i+1}-b_i)} \\
&\geq \Theta(b_{i-1})e^{c_o(a_i-b_{i-1})}e^{c_o(a_{i+1}-b_i)} \\
&= \Theta(b_{i-1})e^{c_o(a_{i+1}-b_i)+(a_i-b_{i-1})}\,.
\end{aligned}$$

Therefore by induction we have

$$\Theta(a_{N+1}) \geq \Theta(b_1)e^{c_o \sum_{i=1}^{N}(a_{i+1}-b_i)} . \tag{4.10}$$

For $r = a_{N+1}$ we have $\alpha'(a_N) \leq 1+\eta$ and $\alpha(a_N) \leq a_N + C$, therefore

$$\Theta(a_N) \leq c_1$$

as in the proof of lemma 4.4, where c_1 is a positive constant independent of N. This is a contradiction, as we have that the set $U_{i=1}^{\beta}[b_i, a_{i+1}]$ has infinite measure. Therefore there is a point r_c such that $\alpha(r_c) > r_c + C$. Lemma 4.1 shows that $\alpha(r) > r + C$ for all $r > r_c$. **Q.E.D.**

Theorem 4.11. *For $p > 2$, let $F(r,\varphi) = (\alpha(r), \varphi)$ be a rotationally symmetric p-harmonic map from the hyperbolic space to itself, where $\alpha \in C^2(0,\infty)$ and $\lim_{r\to 0^+}\alpha(r) = \alpha(0) = 0$. Assume that there exists $r_o > (\ln 3)/2$ such that $\alpha(r_o) < r_o$ and $\alpha'(r_o) < 1$. Then α is bounded on $[0,\infty)$.*

Proof. It follows from lemma 4.1 that there is a positive constant $\epsilon_o > 0$ such that $\alpha(r) \leq r - \epsilon_o$ for all $r > r_o$. At a point $r > r_o$, let $\alpha(r) = r - \epsilon$ where $\epsilon \geq \epsilon_o$. We have

$$\begin{aligned}
& \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
= \; & \frac{2g(\alpha(r))g'(\alpha(r))}{f^{\frac{3}{2}}(r)f^{\prime\frac{1}{2}}(r)}[\frac{f'(r)(\alpha'(r))^2}{f(r)}]^{\frac{1}{2}} - \frac{f'(r)g^2(\alpha(r))}{f^3(r)} - \frac{f'(r)}{f(r)}(\alpha'(r))^2 \\
\leq \; & \frac{1}{1-\tau}\frac{g^2(\alpha(r))g'^2(\alpha(r))}{f^3(r)f'(r)} + (1-\tau)\frac{f'(r)}{f(r)}(\alpha'(r))^2 - \frac{f'(r)g^2(\alpha(r))}{f^3(r)} \\
& \quad -\frac{f'(r)}{f(r)}(\alpha'(r))^2 \\
= \; & \frac{1}{1-\tau}\frac{g^2(\alpha(r))g'^2(\alpha(r))}{f^3(r)f'(r)} - (1-\tau)\frac{f'(r)g^2(\alpha(r))}{f^3(r)} \\
& \quad -\tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \\
= \; & \frac{g^2(\alpha(r))}{4f^3(r)f'(r)}[\frac{e^{2\alpha(r)} + 2 + e^{-2\alpha(r)}}{1-\tau} - (1-\tau)(e^{2r} + 2 + e^{-2r})] \\
& \quad -\tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \\
= \; & \frac{g^2(\alpha(r))}{4f^3(r)f'(r)}(\frac{e^{2r}e^{-2\epsilon} + 2 + e^{-2r}e^{2\epsilon} - (1-\tau)^2(e^{2r} + 2 + e^{-2r})}{1-\tau}) \\
& \quad -\tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2]
\end{aligned}$$

$$\begin{aligned}
&= -\frac{g^2(\alpha(r))e^{2r}}{4f^3(r)f'(r)}[\frac{(1-\tau)^2 - e^{-2\epsilon} - 2e^{-2r}[(1-\tau)^2+1] - e^{-4r}(e^{2\epsilon} - (1-\tau)^2)}{1-\tau}] \\
&\qquad -\tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \\
&\le -\frac{g^2(\alpha(r))e^{2r}}{4f^3(r)f'(r)}[\frac{(1-\tau)^2 - e^{-2\epsilon_o} - 2e^{-2r}[(1-\tau)^2+1] - e^{-2r} + e^{-4r}(1-\tau)^2}{1-\tau}] \\
&\qquad -\tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2]\,,
\end{aligned}$$

where we have made use of the inequality $2AB \le A^2/(1-\tau) + (1-\tau)B^2$ for $1 > \tau > 0$, and $r \ge \epsilon \ge \epsilon_o > 0$. Choose $\tau > 0$ such that $(1-\tau)^2 \ge e^{-2\epsilon_o} + (1-e^{-2\epsilon_o})/2 = (1+e^{-2\epsilon_o})/2$. Since $g'(r) = f'(r) = (e^r + e^{-r})/2$, we can find positive constants $\bar{r}$, such that for $r > \bar{r}$, we have

(4.12)
$$\frac{g^2(\alpha(r))e^{2r}}{4f^3(r)f'(r)}[\frac{(1-\tau)^2 - e^{-2\epsilon_o} - 2e^{-2r}[(1-\tau)^2+1] - e^{-2r} + (1-\tau)^2)e^{-4r}}{1-\tau}] > 0\,.$$

Hence for all $r > \bar{r}$,

$$\begin{aligned}
&\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
\le\ & -\tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \\
=\ & -\tau\frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
\le\ & -c[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\,,
\end{aligned}$$

where $c > 0$ is a positive constant, as $f'(r)/f(r)$ is a bounded function for $r > \bar{r} > 0$. Hence

$$\Theta'(r)[(p-2)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha)}{f^2(r)}] \le -c\,[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\Theta(r)\,. \tag{4.13}$$

If we take $b = \max\{p-2, n-1\}$ and $c' = c/b$, then

$$\Theta'(r) \le -c'\Theta(r) \quad \text{for all} \quad r > \bar{r}\,. \tag{4.14}$$

Hence

$$\Theta(r) \le e^{-c'(t-\bar{r})}\Theta(\bar{r})\,.$$

So we have

$$\alpha'(r) \le C'e^{-\frac{c'(t-\bar{r})}{p-2}} \quad \text{for all} \quad r > \bar{r}\,,$$

where $C' = (\Theta(\bar{r}))^{\frac{1}{p-2}}$ is a positive constant unless α is the trivial solution. As $\alpha'(r) \geq 0$, upon integration we have that α is bounded. **Q.E.D.**

Remark. As $\bar{r}$ can be estimated by using ϵ_o and r_o and thus $\Theta(\bar{r})$ can be estimated in terms of ϵ_o and r_o. Therefore we can find a constant $C(r_o, \epsilon_o)$ such that $\alpha(r) \leq C(r_o, \epsilon_o)$ for all $r > 0$. As we have $\alpha'(r) > 0$ for all $r > 0$, if α is not a trivial solution, hence under the assumption of theorem 4.1, there is a positive number a such that $\lim_{r\to\infty} \alpha(r) = a$.

Theorem 4.15. *For $p > 2$, let $F(r, \varphi) = (\alpha(r), \varphi)$ be a rotationally symmetric p-harmonic map from the hyperbolic space to itself, where $\alpha \in C^2(0, \infty)$ and $\lim_{r\to 0^+} \alpha(r) = \alpha(0) = 0$. Suppose that there exist positive constants C and $\bar{r}$ such that $r + C \geq \alpha(r) \geq r - C$ for all $r > \bar{r}$, then α is asymptotic to the the identity map $\alpha_I(r) = r$*

Proof. By lemma 4.4, we can find a constant r_1 such that either $\alpha(r) \geq r$ or $\alpha(r) \leq r$ for all $r > r_1$. Assume that there is a point $r_o > \max\{\bar{r}\,, r_1\}$ such that $\alpha(r_o) > r_o$. If there is a point $r_2 > r_o$ such that $\alpha(r_2) = r_2$, then either $\alpha(r) = r$ for all $r \geq r_2$ or there is a point r_3 such that $\alpha(r_3) > r_3$. In the second case then we can find a point r_4 such that $r_2 < r_4 \leq r_3$ and $\alpha(r_4) > r_4$ and $\alpha'(r_4) > 1$. By theorem 4.5, $\alpha(r) > r + C$ when r is large. Therefore $\alpha(r) > r$ for all $r > r_o$ if $\alpha(r) \not\equiv r$ on (r_2, ∞). Let

$$c_o = \inf\,\{c > 0 \mid \text{there exists a point } r_c > r_o \text{ such that } \alpha(r_c) = r_c + c\}\,.$$

If $\alpha(r_c) = r_c + c$ for some $r_c > r_o$, then $\alpha(r) \leq r + c$ for all $r > r_c$. For if there is a point r_5 such that $r_5 > r_o$ and $\alpha(r_5) > r_5 + c$, then we can find $r_5' \in (r_c\,, c_5]$ such that $\alpha(r_5') > r_5'$ and $\alpha'(r_5) > 1$. By theorem 4.5 this is not possible in this case. Given any $\epsilon > 0$, we have a point $r_{c_o+\epsilon}$ such that $\alpha(r_c) = r_{c_o+\epsilon} + c_o + \epsilon$ and $\alpha(r) \leq r + c_o + \epsilon$ for all $r \geq r_{c_o+\epsilon}$. As $\alpha(r) > r + c_o$ for all $r > r_o$, we have the line $y = x + c_o$ being an asymptotic of α. If $c_o > 0$, then we have $\alpha(r) > r + c_o$ for all $r > r_o$. Theorem 4.5 shows that $\alpha'(r) \leq 1$ for all $r > r_o$. In fact if there is a point $r > r_o$ such that $\alpha'(r) = 1$, then as in (4.2) and the proof of lemma 4.1, we have $\alpha''(r) > 0$. But this is not possible by theorem 4.5. Thus $\alpha'(r) < 1$ for all $r > r_o$. Since $\alpha(r) \geq r + c_o$ for all $r > r_o$, we can find sequence $r_i > r_o$, $i \in \mathbf{N}$ with $\lim_{i\to\infty} r_i = \infty$, such that $\alpha(r_i) > r_i + c_o$ with $c_o > 0$ and $1 - \eta \leq \alpha'(r_i) < 1$, where we can choose η as small as we like. We choose $\eta > 0$ so small such that $2e^{2c_o}(1 - \eta) > 2 + (e^{2\epsilon_o} - 1)/2$, that is, $3e^{2\epsilon_o} - 4\eta > 3$. Then for any point r with $\alpha'(r) \geq 1 - \eta$ and $\alpha(r) = r + c_o'$ with $C \geq c_o' \geq c_o$, a calculation as in (4.2) shows that we can find a constant $r_6 > r_o$ such that if $r > r_6$,

$$\begin{aligned}(4.16)\qquad & \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\ \geq\ & \frac{1}{8f^3(r)}[2(1-\eta)(e^{3r}e^{2c_o} - e^{-r}e^{-2c_o} - e^r e^{2c'_o} + e^{-3r}e^{-2c'_o}) \\ & -(e^{3r}e^{2c'_o} - 2e^r + e^{-r}e^{-2c'_o} + e^r e^{2c'_o} - 2e^{-r} + e^{-3r}e^{-2c'_o}) \\ & -(e^{3r} - e^r - e^{-r} + e^{-3r})] \\ \geq\ & \frac{1}{8f^3(r)}[C'(C,\eta,c_o)e^{3r}]\end{aligned}$$

where $C'(C,\eta,c_o)$ is a positive constant. As $f(r) = (e^r + e^{-r})/2$, therefore we can find a point $r_8 > r_7$ such that

$$(4.17)\qquad \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \geq C''(\eta,c_o)$$

for $r > r_7$ and for some constant $C''(\eta,c_o) > 0$. That is,

$$\Theta'(r)[(p-2)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \geq (p-2)(n-1)C''(\eta,c_o)\Theta(r)$$

for all $r > r_8$. Since $1-\eta < (\alpha'(r))^2 < 1$ and $C_o \leq g^2(\alpha(r))/f^2(r)) \leq C'_o$, as $r < \alpha(r) \leq r + C$ for all $r > r_8$, we have $C_1 < \theta(r) < C'_1$ for all $r > r_8$ and for some positive constant C_o, C'_o, C_1 and C'_1. Furthermore $(\frac{g^2(\alpha)}{f^2})' < 0$ as in the proof of lemma 4.1. Therefore $\alpha''(r) \geq c_1$ if $r > r_7$ and $\alpha'(r) \geq 1-\eta$, where c_1 is a positive constant. For the sequence $\{x_i\}$, we choose a point $x_i > r_7$ and at x_i we have $\alpha(x_i) > x_i + c_o$ and $1 > \alpha(x_i) \geq 1-\eta$. Therefore $\alpha''(x_i) \geq c_1$. And for all $r > x_i$ we have $\alpha(x_i) > x_i + c_o$ and $\alpha(x_i) \geq 1-\eta$ as $\alpha''(r) \geq c_1 > 0$ for all $r \geq r_i$. Thus $\alpha''(r) \geq c_1$ for all $r \geq r_i$. Then we can find $r_8 > r_i$ such that $\alpha(r_8) = 1$, which is impossible. Hence we must have $c_o = 0$. Similar argument works for $\alpha(r) < r$, using theorem 4.11. **Q.E.D.**

For $p > 2$, let $F(r,\varphi) = (\alpha(r),\varphi)$ be a rotationally symmetric p-harmonic map from the hyperbolic space to itself, where $\alpha \in C^2(0,\infty)$ and $\lim_{r\to 0^+}\alpha(r) = \alpha(0) = 0$. By lemma 4.3, there is a point $r_o > 0$ such that either $\alpha(r) \geq r$ or $\alpha(r) \leq r$ for all $r > r_o$. If $\alpha(r) \not\equiv r$ for $r > r_o$, then there is a point $r_i > r_o$ such that either $\alpha(r_i) > r_i$ or $\alpha(r_i) < r_i$. If $\alpha(r_i) > r_i$ and $\alpha'(r_i) > 1$, then by theorem 4.4, $\alpha(r) > r + C$ for all r large. If $\alpha(r_i) < r_i$ and $\alpha'(r_1) < 1$, then α is bounded. The other cases are $\alpha'(r) \leq 1$ for all $r \geq r_1$ with $\alpha(r) > r$ and $\alpha'(r) \geq 1$ for all $r \geq r_1$ with $\alpha(r) < r$, that is, $\alpha(r)$ is bounded between $r + C$ and $r - C$ for some $C > 0$ and for all r large. Hence α is asymptotic to the line $y = x$ by theorem

4.15. Therefore we have the following.

Theorem 4.18. *For $p > 2$, let $F(r, \varphi) = (\alpha(r), \varphi)$ be a rotationally symmetric p-harmonic map from the hyperbolic space to itself, where $\alpha \in C^2(0, \infty)$ and* $\lim_{r\to 0^+} \alpha(r) = \alpha(0) = 0$. *Then either 1) for any constant $C > 0$, $\alpha(r) > r + C$ for all r large, or 2) α is bounded, or 3) α is asymptotic to the identity map.*

5. Asymptotic properties

Let f, g be functions in $C^2([0, \infty))$ which satisfy the conditions in (1.1). We study asymptotic properties of solutions to (1.4). In this section we assume that f is similar to the hyperbolic space and g is similar to the Euclidean space. The result in this section can be founded in [9] We assume that there are constants $a, C > 0$ such that

$$\frac{1}{C}e^{ar} \le f(r) \le Ce^{ar}\,, \qquad \frac{1}{C}e^{ar} \le f'(r) \le Ce^{ar} \tag{A}$$

for all $r > 1$. And for $y > 1$, we assume that

$$g(y) \le C'y^m\,, \qquad 0 \le g'(y) \le C'y^{m-1} \tag{B}$$

for some constants $m \ge 1$ and $C' > 0$.

Lemma 5.1. *For $p > 2$, let $\alpha(r) \in C^2(0, \infty)$ be a positive solution to (1.4) with* $\lim_{r\to 0^+} \alpha(r) = 0$. *Suppose that f and g satisfy the conditions (A) and (B), respectively. For $m > 1$. let $c < a/(m - 1)$ be a positive constant. Then there exists a positive constant r_o such that either $\alpha(r) \ge e^{cr}$ or $\alpha(r) \le e^{cr}$ for all $r > r_o$.*

Proof. If α is bounded, then we have $\alpha(r) \le e^{cr}$ for r large enough. Therefore we may assume that $\lim_{r\to\infty} \alpha(r) = \infty$. Hence there exists a constant $r' > 1$ such that $\alpha(r) > 1$ for all $r > r'$. For $m > 1$, suppose that we cannot find such a r_o as in the statement of the theorem, then for any $r_o > r'$, there exist two points $r_2 > r_1 > r_o$ such that $\alpha(r_1) = e^{cr_1}$, $\alpha(r_2) = e^{cr_2}$ and $\alpha(r') \le e^{cr'}$ for $r' \in [r_1, r_2]$. There is a point $r \in [r_1, r_2]$ such that $\alpha(r) \le e^{cr}$, $\alpha'(r) = ce^{cr}$ and $\alpha''(r) \ge c^2e^{cr}$. If $c < a/(m - 1)$, we have

$$\begin{aligned} &\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\ \le\ &\frac{2C'^2C^2ce^{2mcr}}{e^{2ar}} - \frac{c^2e^{2cr}}{C^2} < -c' \end{aligned} \tag{5.2}$$

if r_o is large enough. Here c' is a positive constant. By (2.3) and (5.2) we have

$$\Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \le -c'(p-2)(n-1)\Theta(r)\,. \tag{5.3}$$

And

$$\begin{aligned} & \Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \\ = \;& (\frac{p}{2}-1)\theta^{\frac{p}{2}-2}(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \\ & \times[(n-1)(\frac{g^2(\alpha(r))}{f^2(r)})' + 2\alpha'(r)\alpha''(r)] \\ \ge \;& (\frac{p}{2}-1)\Theta(r)[(n-1)(\frac{g^2(\alpha(r))}{f^2(r)})' + 2\alpha'(r)\alpha''(r)]\,, \end{aligned} \tag{5.4}$$

as $\theta(r) = (n-1)\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2$ and $\Theta(r) = \theta^{\frac{p}{2}-1}(r)$. By (5.3), there is a constant $c'' > 0$ such that at r we have

$$(n-1)(\frac{g^2(\alpha(r))}{f^2(r)})' + 2\alpha'(r)\alpha''(r) \le -c''\,. \tag{5.5}$$

On the other hand

$$\begin{aligned} (\frac{g^2(\alpha)}{f^2(r)})' &= \frac{2f(r)g(\alpha)[f(r)g'(\alpha)\alpha'(r) - f'(r)g(\alpha)]}{f^4(r)} \\ &\ge -\frac{2f'(r)g^2(\alpha)}{f^3(r)} \\ &\ge -2C''^2C^2\frac{e^{2mcr}}{e^{2ar}} \\ &\ge -\epsilon e^{2cr}\,, \end{aligned} \tag{5.6}$$

as $c < a/(m-1)$. Here ϵ is a positive constant. And by choosing r_o large, we may assume that $\epsilon < 2c^3/(n-1)$. There if r_o is large and $c < a/(m-1)$, then (5.5) and (5.6) imply that

$$2\alpha'(r)\alpha''(r) \le (n-1)\epsilon c^{2cr}\,.$$

That is

$$\alpha''(r) \le \frac{(n-1)\epsilon}{2c}c^{cr} < c^2 e^{cr}\,,$$

which is a contradiction. Hence there exists a positive constant $r_o > r'$ such that either $\alpha(r) \ge e^{cr}$ or $\alpha(r) \le e^{cr}$ for all $r > r_o$. **Q.E.D.**

Lemma 5.7. *For $p > 2$, let $\alpha(r) \in C^2(0,\infty)$ be a positive solution to (1.4) with $\lim_{r\to 0^+}\alpha(r) = 0$. Suppose that f and g satisfy the conditions (A) and (B)* *with $m > 1$, respectively. Let $c < a/(m-1)$ be a positive constant. Suppose that there exists a $r_o > 0$ such that $\alpha(r) \le e^{cr}$ for all $r > r_o$, then $\alpha(r) \le C$ for some positive constant C.*

Proof. Assume that $\alpha(r) \le e^{cr}$ for all $r > r_o$, where $c < a/(m-1)$. We may assume that $\alpha \not\equiv 0$. We can find positive constants r' and δ such that $\alpha(r) \ge \delta$ for all $r \ge r'$. There is a positive constant C'' such that

$$g(y) \le C'' y^m\,, \qquad 0 \le g'(y) \le C'' y^{m-1}$$

for all $y \ge \delta$. We may take $r_o > r' > 1$. Then for $r > r_o$ and $1 > \tau > 0$, we have

$$\begin{aligned}
(5.8)\qquad & \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
= \quad & \frac{2g(\alpha(r))g'(\alpha(r))}{f^{\frac{3}{2}}(r)f'^{\frac{1}{2}}(r)}[\frac{f'(r)(\alpha'(r))^2}{f(r)}]^{\frac{1}{2}} - \frac{f'(r)g^2(\alpha(r))}{f^3(r)} - \frac{f'(r)}{f(r)}(\alpha'(r))^2 \\
\le \quad & \frac{1}{1-\tau}\frac{g^2(\alpha(r))g'^2(\alpha(r))}{f^3(r)f'(r)} + (1-\tau)\frac{f'(r)}{f(r)}(\alpha'(r))^2 - \frac{f'(r)g^2(\alpha(r))}{f^3(r)} - \frac{f'(r)}{f(r)}(\alpha'(r))^2 \\
= \quad & \frac{g^2(\alpha(r))}{(1-\tau)f^3(r)f'(r)}[g'^2(\alpha(r)) - (1-\tau)^2 f'(r)^2] - \tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \\
\le \quad & \frac{g^2(\alpha(r))}{(1-\tau)f^3(r)f'(r)}[C''^2 e^{2c(m-1)r} - \frac{(1-\tau)^2}{C^2}e^{2ar}] - \tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2]\,,
\end{aligned}$$

where we have used the inequality $2AB \le (1-\tau)A^2 + 1/(1-\tau)B^2$ for $0 < \tau < 1$. Hence there exists a constant $r'' > r_o$ such that for $r > r''$ we have

$$C''^2 e^{2c(m-1)r} - \frac{(1-\tau)^2}{C^2}e^{2ar} \le 0\,.$$

For $r > r''$ we have

$$\begin{aligned}
(5.9)\qquad & \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
= \quad & -\tau\frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
\le \quad & \frac{-\tau}{C^2}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\,.
\end{aligned}$$

Together with (2.3) we have

$$\begin{aligned}
(5.10)\qquad & \Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha)}{f^2(r)}] \\
\le \quad & -\frac{\tau}{C^2}(p-2)(n-1)\,[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\Theta(r)\,.
\end{aligned}$$

Thus we have

$$(5.11) \qquad \Theta'(r) \leq -c'\Theta(r) \quad \text{for all} \quad r > r'' \,,$$

where $c' = (p-2)(n-1)\tau/(bC^2)$ and $b = \max\{p-2, n-1\}$ are positive constants. Integrating both sides of the above inequality we have

$$\Theta(r) \leq e^{-c'(t-\bar{r})}\Theta(r'')\,, \quad r > r''\,.$$

So we have

$$\alpha'(r) \leq C_1 e^{-\frac{c'(t-r'')}{p-2}} \quad \text{for all} \quad r > r''\,,$$

where $C_1 = (\Theta(r''))^{\frac{1}{p-2}}$ is a positive constant. As $\alpha'(r) > 0$, upon integration we conclude that α is bounded. **Q.E.D.**

Combining the above two lemmas we have the following.

Theorem 5.12. *For $p > 2$, let $\alpha(r) \in C^2(0,\infty)$ be a positive solution to (1.4) with $\lim_{r\to 0^+}\alpha(r) = 0$. Suppose that f and g satisfy the conditions (A) and (B) with $m > 1$, respectively. Let $c < a/(m-1)$ be a positive constant. Then either there exists a positive constant r_o such that $\alpha(r) \geq e^{cr}$ for all $r > r_o$ or there is a finite positive number C such that $\alpha(r) \leq C$ for all $r > 0$.*

In the following we assume that

$$\text{(C)} \qquad g'(y) \leq C_2 g(y) \qquad \text{for all} \quad y \geq 1\,,$$

where C_2 is a positive constant. Condition (C) implies that g grows at most exponentially. Condition (B) is stronger than condition (C). The following could be considered as a generalization of lemma 5.7.

Lemma 5.13. *For $p > 2$, let $\alpha(r) \in C^2(0,\infty)$ be a positive solution to (1.4) with $\lim_{r\to 0^+}\alpha(r) = 0$. Suppose that f and g satisfy the conditions (A) and (C), respectively, and $g'(y) > 0$ for all $y > 0$. There is a positive constant $c > 0$ such that if there exists a positive constant r_o such that $\alpha(r) \leq g^{-1}(cf(r))$ for all $r > r_o$, then α is a bounded function on $\mathbb{R}^+$.*

Proof. We can find positive constants r' and δ such that $\alpha(r) \geq \delta$ for all $r \geq r'$. There is a positive constant C_3 such that

$$g'(y) \leq C_3 g(y) \qquad \text{for all} \quad y \geq \delta\,.$$

For $0 < \tau < 1$ and $r > r'$, from (5.8) we have

$$\begin{aligned} & \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\ \leq \; & \frac{g^2(\alpha(r))}{(1-\tau)f^3(r)f'(r)}[g'^2(\alpha(r)) - (1-\tau)^2 f'(r)^2] - \tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \\ \leq \; & \frac{g^2(\alpha(r))}{(1-\tau)f^3(r)f'(r)}[C_3^2 g^2(\alpha(r)) - \frac{(1-\tau)^2}{C^2} f^2(r)] - \tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \, . \end{aligned}$$

If we choose a positive constant c such that $c \leq (1-\tau)/(CC_3)$, and if $\alpha(r) \leq g^{-1}(cf(r))$, then we have

$$\frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \leq -\tau[\frac{f'(r)g^2(\alpha(r))}{f^3(r)} + \frac{f'(r)}{f(r)}(\alpha'(r))^2] \, .$$

We can process as in the proof of lemma 5.7. **Q.E.D.**

The above lemma provides asymptotic information about $\alpha(r)$. We can also obtain medium range information about $\alpha(r)$.

Lemma 5.14 *For $p > 2$, let $\alpha(r) \in C^2(0, \infty)$ be a positive solution to (1.4) with $\lim_{r \to 0^+} \alpha(r) = 0$. Suppose that f and g satisfy the conditions (A) and (C), respectively, and $g'(y) \geq 0$ for all $y > 0$. Assume that α is not a bounded function on $\mathbf{R}^+$ and let $\alpha(\bar{r}) = 1$ for some $\bar{r} > 0$. Then there exists a positive constant δ such that the energy density $\theta(r) \geq \delta$ for all $r \geq \bar{r}$.*

Proof. From (2.3) we have

$$\begin{aligned} & \Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \\ = \; & (p-2)\Theta(r)\{\frac{2(n-1)g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{1}{2}\frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\ & -(n-\frac{3}{2})\frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\} \, . \end{aligned} \tag{5.15}$$

At a point $r \geq \bar{r}$, we have $\alpha(r) \geq 1$. So at the point $r > \bar{r}$, we have either

(i) $$\frac{2(n-1)g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} \geq \frac{1}{2}\frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \, ,$$

or

(ii) $$\begin{aligned} & \Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \\ \leq \; & -(p-2)(n-\frac{3}{2})\Theta(r)\frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \, . \end{aligned}$$

In case of (i), from condition (A) and (C) we have

$$2C_2(n-1)\alpha'(r) \geq \frac{f'(r)f(r)(\alpha'(r))^2}{g^2(\alpha(r))} + \frac{f'(r)}{2f(r)} \geq \frac{1}{2C^2}\,.$$

By choosing a bigger constant if necessary, we may assume that (A) holds for $r > \bar{r}$. Thus if (i) holds at $r \geq \bar{r}$, we have

$$\alpha'(r) \geq \frac{1}{4(n-1)C_2C^2}\,. \tag{5.16}$$

Take

$$\delta = (\frac{1}{8(n-1)C_2C^2})^2\,. \tag{5.17}$$

Suppose that there is a point $r'' \geq r_o$ such that $\theta(r'') \leq \delta$. Let

$$O = \{t \in [r'', \infty) \mid \theta(r) \leq \delta \ \text{ for all } \ r \in [r'', t]\}\,.$$

Therefore $r' \in 0$ and

$$\alpha'(r') \leq \frac{1}{8(n-1)C_2C^2}\,.$$

We can find t such that $t > r'$ and

$$\alpha'(r) \leq \frac{1}{4(n-1)C''C^2}$$

for all $r \in [r'', t]$. Hence we have (ii) for all $r \in [r'', t]$. Thus $\Theta'(r) \leq 0$ and hence $\theta'(r) \leq 0$ for all $r \in [r'', t]$. We have $\theta(r) \leq \delta$ for all $r \in [r'', t]$. Suppose that sup $O = a < \infty$. Then at the point a we have $\theta(a) \leq \delta$. The same argument shows that there exists a positive number ϵ such that $\Theta'(r) \leq 0$ for all $r \in (a, a+\epsilon)$. Then $a + \epsilon \in O$, which is a contradiction. Therefore we have sup $O = \infty$, or

$$\theta(r) \leq \delta \ \text{ for all } \ r \in [r', \infty)\,. \tag{5.18}$$

In particular,

$$\alpha'(r) < \frac{1}{4(n-1)C''C^2}$$

for all $r \in [r', \infty)$. By (ii) we have

$$\Theta'(r)[(p-1)(\alpha'(r))^2 + (n-1)\frac{g^2(\alpha(r))}{f^2(r)}] \leq -\frac{(p-2)(n-\frac{3}{2})}{C^2}\Theta[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2]\,.$$

A similar argument as in the proof of lemma 5.7 shows that there is a positive constant $c > 0$ such that $\alpha(r) \leq c$ for all $r \in (0, \infty)$. This contradicts the assumption that α is not bounded. Therefore $\theta(r) \geq \delta$ for all $r \geq \bar{r}$. **Q.E.D.**

Theorem 5.19 *For $p > 2$, let $\alpha(r) \in C^2(0,\infty)$ be a positive solution to (1.4) with $\lim_{r\to 0^+} \alpha(r) = 0$. Suppose that f and g satisfy the conditions (A) and (C), respectively, and $g'(y) > 0$ for all $y > 0$. Then either there exist positive constants r_o and c_o such that either $\alpha(r) \geq c_o(r - r_o)$ for all $r > r_o$, or α is a bounded function on $\mathbf{R}^+$.*

Proof. If α is not bounded, then there is a positive constant $\bar{r}$ such that $\alpha(\bar{r}) = 1$. By lemma 5.14, we have $\theta(r) \geq \delta$ for all $r \geq \bar{r}$. With the conditions (A) and (C), we can find positive constants c' and r'_o such that

$$g^{-1}(\frac{\delta}{\sqrt{2(n-1)}} f(r)) \geq c' r$$

for all $r > r'_o$. We can choose $c_o < \min\{c', \sqrt{\delta/2}\}$ and $r_o > \max\{\bar{r}\,, r'_o\}$. If there is a point $r' > r_o$ such that $\alpha(r') < c_o(r' - r_o)$. Then we can find a point $r > r_o$ such that $\alpha(r) < c_o r$ and $\alpha'(r) < c_o$. Thus

$$0 < \alpha'(r') < \sqrt{\frac{\delta}{2}}\,.$$

And

$$(n-1)\frac{g^2(\alpha(r'))}{f^2(r)} \leq (n-1)\frac{g^2(cr)}{f^2(r)} \leq (n-1)\frac{g^2(g^{-1}(\frac{\delta}{\sqrt{2(n-1)}} f(r)))}{f^2(r)} \leq \frac{\delta}{2}\,,$$

as $g' \geq 0$ implies that g is non-decreasing. Therefore $\theta < \delta$, contradiction. **Q.E.D.**

6. Metrics with polynomial growth

For $p > 2$, let $F(r,\varphi) = (\alpha(r), \varphi)$ be a rotationally symmetric p-harmonic map from $M^n(f)$ to itself. The results in this section can be founded in [8].

Lemma 6.1. *For $f(r) = r^m$ with $m \geq 1$, if there exists a point $r_o > 0$ such that $\alpha(r_o) < cr$ and $\alpha'(r_o) < c$ for some $c \in (0,1]$, then $\alpha(r) < cr$ for all $r \geq r_o$.*

Proof. Assume that there is a point $r' > r_o$ such that $\alpha(r') = cr'$. As $\alpha'(r_o) < c$, there exists a point $r \in (r_o, r_1]$ such that $\alpha(r) = c'r$ for some constant $c' \in (0, c)$ and $\alpha'(r) = c$ and $\alpha''(r) \geq 0$. At r, we have

$$
\begin{aligned}
(6.2) \qquad & \frac{2g(\alpha(r))g'(\alpha(r))\alpha'(r)}{f^2(r)} - \frac{f'(r)}{f(r)}[\frac{g^2(\alpha(r))}{f^2(r)} + (\alpha'(r))^2] \\
= \quad & \frac{2m\alpha^{2m-1}c}{r^{2m}} - \frac{m}{r}(\frac{\alpha^{2m}}{r^{2m}} + c^2) \\
= \quad & \frac{m}{r}(2cc'^{2m-1} - c'^{2m} - c^2)
\end{aligned}
$$

It can be shown that if $m \geq 1$, then $2cc'^{2m-1} - c'^{2m} - c^2 < 0$ for $c' \in (0, c)$. In fact, if we let

$$\phi(x) = 2cx^{2m-1} - x^{2m} - c^2\,,$$

then $\phi(c) = c^{2m} - c^2 \leq 0$ as $c \leq 1$, $m \geq 1$, and $\phi'(x) > 0$ for $x \in (0, c)$. Hence $\Theta'(r) < 0$. Therefore

$$0 > \theta'(r) = (\frac{g^2(\alpha)}{f^2(r)})' + 2\alpha'(r)\alpha''(r)\,.$$

And

$$(6.3) \qquad (\frac{g^2(\alpha)}{f^2(r)})' = \frac{2m}{r^{4m}}(r^{2m}\alpha^{2m-1}c - \alpha^{2m}r^{2m-1}) = \frac{2m}{r}(c'^{2m-1}c - c'^{2m}) > 0$$

as $0 < c' < c$. Hence we have $\alpha''(r) > 0$, contradiction. **Q.E.D.**

Theorem 6.4. *Let $f(r) = g(r) = r^m$ for some $m \geq 1$. Let $\alpha \in C^1[0, \infty) \cap C^2(0, \infty)$ be a solution to the rotationally symmetric p-harmonic equation with $\alpha(0) = 0$. If $\alpha'(0) = c$ for some $c \in (0, 1]$, then $\alpha(r) \leq cr$ for all $r > 0$.*

Proof. For $\alpha'(0) = c < 1$, assume that there is a point $r' > 0$ such that $\alpha(r') > cr'$. Then we can find $c' > c$ such that $\alpha(r') > c'r'$. Since $\alpha'(0) = c < c'$. we can find a point $r_o < r'$ close to zero such that $\alpha(r_o) < c'r_o$ and $\alpha'(r_o) < c'$. Lemma 6.1 implies that $\alpha(r) < c'r$ for all $r > r_o$, contradiction. So we have $\alpha(r) \leq cr$ for all $r > 0$. In case $c = 1$, $\alpha(r) = r$ is the unique solution to the equation with $\alpha'(0) = 1$ [2]. **Q.E.D.**

For $m = 1$, that is, the Euclidean space, and for any positive number c, the function $\alpha_c = cr$ is a rotationally symmetric p-harmonic map into the Euclidean space for all $p > 0$.

Theorem 6.5. *Let $f(r) = r^m$ with $m > 1$. For $p > 2$, let $F(r, \varphi) = (\alpha(r), \varphi)$ be a rotationally symmetric p-harmonic map from $M^n(f)$ to itself, where $\alpha \in C^2(0, \infty)$*

and $\lim_{r\to 0^+}\alpha(r)=\alpha(0)=0$. *Then for any* $c>0$, *we can find a positive number* r_c *such that either* $\alpha(r)\geq cr$ *or* $\alpha(r)\leq cr$ *for all* $r>r_c$.

Proof. Let $c\in(0,1]$ and suppose that there exist two points $r_2>r_1>0$ such that $\alpha(r_1)=cr_1$, $\alpha(r_2)=cr_2$ and $\alpha(r)\leq cr$ for all $r\in[r_1,r_2]$ but $\alpha(r)\not\equiv cr$ on $[r_1,r_2]$. Then there exist a point $r_o\in(r_1,r_2)$ such that $\alpha(r_o)<cr_o, \alpha'(r_o)<c$. Lemma 6.1 implies $\alpha(r)<cr$ for all $r\geq r_o$, which is a contradiction. Hence we can find a positive number r_c such that either $\alpha(r)\geq cr$ or $\alpha(r)\leq cr$ for all $r>r_c$. Let $c>1$ and suppose that there exist two points $r_2>r_1>0$ such that $\alpha(r_1)=cr_1$, $\alpha(r_2)=cr_2$ and $\alpha(r)\geq cr$ for all $r\in[r_1,r_2]$ but $\alpha(r)\not\equiv cr$ on $[r_1,r_2]$. Let

$$c_o=\inf\{\gamma\geq c\mid \gamma r\geq\alpha(r)\ \text{ for all }\ r\in[r_1,r_2]\}\,.$$

Thus $c_or\geq\alpha(r)$ for all $r\in[r_1,r_2]$. Let

$$c'=\max\{c\,,\frac{m}{2m-1}c_o\}\,.$$

As $m>1$ and $c_o>c$, we have $c'<c_o$. Hence there exist x_1',x_2' such that $x_1\leq x_1'<x_2'\leq x_2$ and $\alpha(r_1)=c'r_1$, $\alpha(r_2)=c'r_2$ and $\alpha(r)\geq c'r$ for all $r\in[r_1',r_2']$. Then there exists a point $r_o\in[r_1',r_2']$ such that $\alpha(r_o)\geq c'r_o, \alpha'(r_o)=c'$ and $\alpha''(r_o)\leq 0$. Let $\alpha(r_o)=c''r_o$ for some $c''\geq c'$. As in (6.2) we have

$$\begin{aligned}&\frac{2g(\alpha(r_o))g'(\alpha(r_o))\alpha'(r_o)}{f^2(r_o)}-\frac{f'(r_o)}{f(r_o)}[\frac{g^2(\alpha(r_o))}{f^2((r_o))}+(\alpha'(r_o))^2]\\ =\;&\frac{m}{t}(2c'c''^{2m-1}-c''^{2m}-c'^2)\,.\end{aligned}$$

The function $\phi(x)=2c'x^{2m-1}-x^{2m}-c'^2$ has $\phi(c')=c'^{2m}-c'^2>0$, since $c'\geq c>1$ and $m>1$. While

$$\phi'(x)=2mx^{2m-2}(\frac{2m-1}{m}c'-x)=\frac{2mx^{2m-2}}{r_o}(\frac{2m-1}{m}c'r_o-xr_o)\,.$$

As $c'\geq m/(2m-1)c_o$, therefore

$$\frac{2m-1}{m}c'r\geq c_or\geq\alpha(r)\quad\text{ for all }\ r\in[r_1',r_2']\,.$$

In particular

$$\frac{2m-1}{m}c'r_o\geq\alpha(r_o)=c''r_o\geq xr_o\quad\text{for all}\ \ x\in[c',c'']\,.$$

Thus $\phi'(x)\geq 0$ for all $x\in[c',c'']$. Hence $\phi(c'')>0$. Therefore

$$\frac{2g(\alpha(r_o))g'(\alpha(r_o))\alpha'(r_o)}{f^2(r_o)}-\frac{f'(r_o)}{f(r_o)}[\frac{g^2(\alpha(r_o))}{f^2((r_o))}+(\alpha'(r_o))^2]>0\,. \tag{6.6}$$

As in (6.3)

$$(\frac{g^2(\alpha(r_o))}{f^2(r_o)})' = \frac{2m}{r_o^{4m}}(r_o^{2m}\alpha^{2m-1}(r_o)c' - \alpha^{2m}(r_o)r_o^{2m-1}) \le 0\,, \tag{6.7}$$

as $\alpha(r_o) \ge c'r_o$. Hence $\alpha''(r_o) > 0$. This is a contradiction. So we can find a positive number r_c such that either $\alpha(r) \ge cr$ or $\alpha(r) \le cr$ for all $r > r_c$. **Q.E.D.**

ACKNOWLEDGEMENT. We would like to thank Professor J.B. McLeod for valuable discussions.

References

[1] Cheung, L.-F., & Law, C.-K., *An intial value approach to rotationally symmetric harmonic maps,* Preprint, 1995.

[2] Cheung, L.-F., Law, C.-K. & Leung, M.-C., *Entire solutions of quasilinear differential equations cooresponding to p-harmonic maps,* Preprint, 1995.

[3] Coddington, E.A. & Levinson, N., *"Theory of Ordinary Differential Equations"*, (1955) McGraw-Hill.

[4] Eells, J. & Lemaire, L., *Another report of harmonic maps,* Bulletin of the London Math. Soc. **20** (1988), 385-524.

[5] Greene, R. & Wu, H., *Function Theory on Manifolds Which Possess a Pole,* Lectures Notes in Math. vol. 699, Springer-Verlag, New York, 1979.

[6] Heinonen, J., Kilpeläinen, T. & Martio, O., *Nonlinear Potential Theory of Degenerate Elliptic Equations,* Oxford Mathematical Monographs, Oxford University Press, Oxford, 1994.

[7] Leung, M.-C., *On the infinitesmal rigidity of harmonic maps,* Preprint, 1995.

[8] Leung, M.-C., *Asymptotic behavior of rotationally symmetric p-harmonic maps,* to appear.

[9] Leung, M.-C., *Positive solutions of second order quasilinear equations corresponding to p-harmonic maps,* Preprint, 1996.

[10] Nakauchi, N. & Takakuwa, S., *A Remark on p-harmonic maps,* Nonlinear Analysis, Theory, Method & Applications **25** (1995), 169-185.

[11] Ratto, A. & Rigoli, M., *On the asymptotic behaviour of rotationally symmetric harmonic maps*, Journal of Diff. Eqn. **101** (1993) 15-27.

[12] Takakuwa, S., *Stability and Liouville theorems of p-harmonic maps*, Japan J. Math., **17** (1991), 317-332.

[13] Uhlenbeck, K., *Regularity for a class of nonlinear elliptic systems*, Acta Math. **138** (1970) 219-240.

[14] White, B., *Homotopy classes in Sobolev spaces and the existence of energy minimizing maps,* Acta Math. **160** (1988), 1-17.

[15] Xu, X.-W. & Yang, P., *A construction of m-harmonic maps of spheres,* International J. of Math. **4** (1993), 521-533.

[16] Yau, S.-T., *Harmonic functions on complete Riemannian manifolds,* Comm. Pure Appl. Math. **28** (1975), 201-228.